

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 September 2005 (15.09.2005)

PCT

(10) International Publication Number
WO 2005/085251 A1

(51) International Patent Classification⁷: **C07D 487/04**,
519/00, A61K 31/5517, A61P 35/00

(21) International Application Number:
PCT/GB2005/000768

(22) International Filing Date: 1 March 2005 (01.03.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0404575.3 1 March 2004 (01.03.2004) GB
0426392.7 1 December 2004 (01.12.2004) GB

(71) Applicant (for all designated States except US):
SPIROGEN LIMITED [GB/GB]; 79 George Street,
Ryde Isle of Wight PO33 2JF (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HOWARD, Philip**,
Wilson [GB/GB]; 3 Westbourne Mews, The Maltings,
St Albans Hertfordshire AL1 1LS (GB). **GREGSON**,
Stephen, John [GB/GB]; 6 Warepoint Drive, Thames-
mead, London Greater London SE28 0HJ (GB).

(74) Agents: **WATSON, Robert** et al.; Mewburn Ellis LLP,
York House, 23 Kingsway, London Greater London WC2B
6HP (GB).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA,
ZM, ZW.


(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 11-HYDROXY-5H-PYRROLO[2,1-C][1,4]BENZODIAZEPIN-5-ONE DERIVATIVES AS KEY INTERMEDIATES
FOR THE PREPARATION OF C2 SUBSTITUTED PYRROLOBENZODIAZEPINES



(I)



(III)

(57) Abstract: The present inventors
have developed a key intermediate for the
production of C2 substituted PBDs, which
has a leaving group at the C2 position,
a carbamate protecting group at the N10
position and a protected hydroxy group
at the C11 position. In a first aspect, the
present invention comprises a compound
with a the formula (I), wherein : R¹⁰ is a
carbamate-based nitrogen protecting group;
R¹¹ is an oxygen protecting group; and R²
is a labile leaving group. In a further aspect,
the present invention comprises a method
of synthesising a compound of formula (III),
or a solvate, thereof, from a compound of
formula (I) as defined in the first aspect, R¹⁶
is either O-R¹¹, wherein R¹¹ is as defined
in the first aspect, or OH, or R¹⁰ and R¹⁶

together form a double bond between N10 and C11; and R¹⁵ is R. The other substituents are defined in the claims. Further aspects
of the present invention relate to compounds of formula (III) (including solvates thereof when R¹⁰ and R¹⁶ form a double bond
between N10 and C11, and pharmaceutical salts thereof), pharmaceutical compositions comprising these, and their use in the
manufacture of a medicament for the treatment of a proliferative disease.

WO 2005/085251 A1


BEST AVAILABLE COPY

11-HYDROXY-5H-PYRROLO[2,1-C][1,4]BENZODIAZEPIN-5-ONE DERIVATIVES AS KEY
INTERMEDIATES FOR THE PREPARATION OF C2 SUBSTITUTED
PYRROLOBENZODIAZEPINES

The present invention relates to pyrrolobenzodiazepines (PBDs), and
in particular pyrrolobenzodiazepines useful in the synthesis of C2
substituted compounds.


Background to the invention

Some pyrrolobenzodiazepines (PBDs) have the ability to recognise
and bond to specific sequences of DNA; the preferred sequence is
PuGpu. The first PBD antitumour antibiotic, anthramycin, was
discovered in 1965 (Leimgruber, et al., *J. Am. Chem. Soc.*, **87**,
5793-5795 (1965); Leimgruber, et al., *J. Am. Chem. Soc.*, **87**, 5791-
5793 (1965)). Since then, a number of naturally occurring PBDs
have been reported, and over 10 synthetic routes have been
developed to a variety of analogues (Thurston, et al., *Chem. Rev.*
1994, 433-465 (1994)). Family members include abbeymycin
(Hochlowski, et al., *J. Antibiotics*, **40**, 145-148 (1987)),
chicamycin (Konishi, et al., *J. Antibiotics*, **37**, 200-206 (1984)),
DC-81 (Japanese Patent 58-180 487; Thurston, et al., *Chem. Brit.*,
26, 767-772 (1990); Bose, et al., *Tetrahedron*, **48**, 751-758 (1992)),
mazethramycin (Kuminoto, et al., *J. Antibiotics*, **33**, 665-667
(1980)), neothramycins A and B (Takeuchi, et al., *J. Antibiotics*,
29, 93-96 (1976)), porothramycin (Tsunakawa, et al., *J.*
Antibiotics, **41**, 1366-1373 (1988)), prothracarcin (Shimizu, et al.,
J. Antibiotics, **29**, 2492-2503 (1982); Langley and Thurston, *J. Org.*
Chem., **52**, 91-97 (1987)), sibanomicin (DC-102) (Hara, et al., *J.*
Antibiotics, **41**, 702-704 (1988); Itoh, et al., *J. Antibiotics*, **41**,
1281-1284 (1988)), sibiromycin (Leber, et al., *J. Am. Chem. Soc.*,
110, 2992-2993 (1988)) and tomamycin (Arima, et al., *J.*
Antibiotics, **25**, 437-444 (1972)). PBDs are of the general
structure:




They differ in the number, type and position of substituents, in both their aromatic A rings and pyrrolo C rings, and in the degree of saturation of the C ring. In the B-ring there is either an imine (N=C), a carbinolamine (NH-CH(OH)), or a carbinolamine methyl ether (NH-CH(OMe)) at the N10-C11 position which is the electrophilic centre responsible for alkylating DNA. All of the known natural products have an (S)-configuration at the chiral C11a position which provides them with a right-handed twist when viewed from the C ring towards the A ring. This gives them the appropriate three-dimensional shape for isohelicity with the minor groove of B-form DNA, leading to a snug fit at the binding site (Kohn, In *Antibiotics III*. Springer-Verlag, New York, pp. 3-11 (1975); Hurley and Needham-VanDevanter, *Acc. Chem. Res.*, **19**, 230-237 (1986)). Their ability to form an adduct in the minor groove, enables them to interfere with DNA processing, hence their use as antitumour agents.

The present inventors have previously disclosed, in PCT/GB2003/004963, cytotoxic compounds having an aryl group at the C2 position, for example:




The synthesis of these compounds was achieved via the following intermediate:



whose synthesis was described in detail in WO 00/12508. This method involves a reduction as a deprotection step, which can lead to overreduction of the compound which is not desirable. Also, with certain C2 groups, the reduction step does not proceed at all.

The following intermediate has also been disclosed:




but its synthesis has proved difficult and only proceeds in low yield.

Disclosure of the invention

The present inventors have developed a key intermediate for the production of C2 substituted PBDs, which has a leaving group at the C2 position, a carbamate protecting group at the N10 position and a protected hydroxy group at the C11 position.

In a first aspect, the present invention comprises a compound with the formula I:




wherein:

R⁶ and R⁹ are independently selected from H, R, OH, OR, SH, SR, NH₂, NHR, NRR', nitro, Me₃Sn and halo;


where R and R' are independently selected from optionally substituted C₁₋₁₂ alkyl, C₃₋₂₀ heterocyclyl and C₅₋₂₀ aryl groups; R⁷ and R⁸ are independently selected from H, R, OH, OR, SH, SR, NH₂, NHR, NHRR', nitro, Me₃Sn and halo, or the compound is a dimer with each monomer being of formula (I), where the R⁷ groups or R⁸ groups of each monomers form together a dimer bridge having the formula -X-R''-X- linking the monomers, where R'' is a C₃₋₁₂ alkylene group, which chain may be interrupted by one or more heteroatoms, e.g. O, S, NH, and/or aromatic rings, e.g. benzene or pyridine, and each X is independently selected from O, S, or NH; or any pair of adjacent groups from R⁶ to R⁹ together form a group -O-(CH₂)_p-O-, where p is 1 or 2; R¹⁰ is a carbamate-based nitrogen protecting group; R¹¹ is an oxygen protecting group; and R² is a labile leaving group.

In a second aspect, the present invention comprises a method of synthesising a compound of formula I as defined in the first aspect of the invention from a compound of formula IIa:



wherein R⁶, R⁷, R⁸, R⁹, R¹⁰ and R¹¹ are as defined in the first aspect; and R¹² and R¹³ together form =O.

It is preferred that the compound of formula IIa is synthesised from a compound of formula IIb:




wherein R^6 , R^7 , R^8 , R^9 , R^{10} and R^{11} are as defined in the first aspect;

R^{12} is $O-R^{14}$, and R^{13} is H, where R^{14} is an oxygen protecting group which is orthogonal to R^{11} .

5

In a third aspect, the present invention comprises a method of synthesising a compound of formula III:



III

or a solvate thereof, from a compound of formula I as defined in the first aspect, wherein R^6 , R^7 , R^8 , R^9 are as defined in the first aspect;

R^{10} is as defined in the first aspect and R^{16} is either $O-R^{11}$, wherein R^{11} is as defined in the first aspect, or OH, or R^{10} and R^{16} together form a double bond between N10 and C11; and
 R^{15} is R.

15


Further aspects of the present invention relate to novel compounds of formula III (including solvates thereof when R^{10} and R^{16} form a double bond between N10 and C11, and pharmaceutical salts thereof), their use in methods of therapy (particularly in treating proliferative diseases), pharmaceutical compositions comprising these, and their use in the manufacture of a medicament for the treatment of a proliferative disease.

20

25 Definitions

Carbamate-based nitrogen protecting groups

Carbamate-based nitrogen protecting groups are well known in the art, and have the following structure:




wherein R'¹⁰ is R as defined above. A large number of suitable groups are described on pages 503 to 549 of Greene, T.W. and Wuts, G.M., *Protective Groups in Organic Synthesis*, 3rd Edition, John Wiley & Sons, Inc., 1999, which is incorporated herein by reference.


Particularly preferred protecting groups include Troc, Teoc, Fmoc, BOC, Doc, Hoc, TcBOC, 1-Adoc and 2-Adoc.

Those protecting groups which can be removed with palladium catalysis are not preferred, e.g. Alloc.


Also suitable for use in the present invention are nitrogen protecting group which can be removed *in vivo* (e.g. enzymatically, using light) as described in WO 00/12507, which is incorporated herein by reference. Examples of these protecting groups include:



, which is nitroreductase labile (e.g. using ADEPT/GDEPT);



and c1ccccc1CC(=O)C*, which are photolabile; and



which is glutathione labile (e.g. using NPEPT).

Oxygen protecting groups

Oxygen protecting groups are well known in the art. A large number of suitable groups are described on pages 23 to 200 of Greene, T.W. and Wuts, G.M., *Protective Groups in Organic Synthesis*, 3rd Edition,

John Wiley & Sons, Inc., 1999, which is incorporated herein by reference.

5 Classes of particular interest include silyl ethers, methyl ethers, alkyl ethers, benzyl ethers, esters, benzoates, carbonates, and sulfonates.

10 Preferred oxygen protecting groups include TBS, THP for the C11 oxygen atom, and methyl ester for the C2 oxygen atom (where present).

15 As mentioned above the oxygen protecting group R^{14} should be orthogonal to the oxygen protecting group R^{11} . Protecting groups which are orthogonal to one another may each be removed using reagents or conditions which do not remove the other protecting group.

20 It may also be preferred that any protecting groups used during the synthesis and use of compounds of formula I are orthogonal to one another. However, it is often not necessary, but may be desirable, for the carbamate-based nitrogen protecting group and R^{11} to be orthogonal to one another, depending on whether the compound of formula III is to be used with the nitrogen protecting group in place.

25

Labile leaving groups

30 Labile leaving groups suitable for use in the present invention are in particular those amenable to palladium catalysed coupling, for example using Suzuki or Stille coupling. Suitable groups include mesylate ($-\text{OSO}_2\text{CH}_3$), $-\text{OSO}_2(\text{C}_n\text{F}_{2n+1})$ where $n=0,1$ or 4 , $-\text{OSO}_2-\text{R}^5$ where R^5 is an optionally substituted phenyl group (e.g. 4-Me-Ph, tosylate), I, Br and Cl. More preferred are $-\text{OSO}_2(\text{C}_n\text{F}_{2n+1})$ where $n=0,1$ or 4 , I, Br and Cl, with triflate ($-\text{OSO}_2\text{CF}_3$) and Br being the most preferred.

35

Substituents

The phrase "optionally substituted" as used herein, pertains to a parent group which may be unsubstituted or which may be substituted.

5

Unless otherwise specified, the term "substituted" as used herein, pertains to a parent group which bears one or more substituents. The term "substituent" is used herein in the conventional sense and refers to a chemical moiety which is covalently attached to, or if appropriate, fused to, a parent group. A wide variety of substituents are well known, and methods for their formation and introduction into a variety of parent groups are also well known.

10

Examples of substituents are described in more detail below.

15

C₁₋₁₂ alkyl: The term "C₁₋₁₂ alkyl" as used herein, pertains to a monovalent moiety obtained by removing a hydrogen atom from a carbon atom of a hydrocarbon compound having from 1 to 12 carbon atoms, which may be aliphatic or alicyclic, and which may be saturated or unsaturated (e.g. partially unsaturated, fully unsaturated). Thus, the term "alkyl" includes the sub-classes alkenyl, alkynyl, cycloalkyl, etc., discussed below.

20

Examples of saturated alkyl groups include, but are not limited to, methyl (C₁), ethyl (C₂), propyl (C₃), butyl (C₄), pentyl (C₅), hexyl (C₆) and heptyl (C₇).

25

Examples of saturated linear alkyl groups include, but are not limited to, methyl (C₁), ethyl (C₂), n-propyl (C₃), n-butyl (C₄), n-pentyl (amyl) (C₅), n-hexyl (C₆) and n-heptyl (C₇).

30

Examples of saturated branched alkyl groups include iso-propyl (C₃), iso-butyl (C₄), sec-butyl (C₄), tert-butyl (C₄), iso-pentyl (C₅), and neo-pentyl (C₅).

35

C₂₋₁₂ Alkenyl: The term "C₂₋₁₂ alkenyl" as used herein, pertains to an alkyl group having one or more carbon-carbon double bonds.

5 Examples of unsaturated alkenyl groups include, but are not limited to, ethenyl (vinyl, -CH=CH₂), 1-propenyl (-CH=CH-CH₃), 2-propenyl (allyl, -CH=CH-CH₂), isopropenyl (1-methylvinyl, -C(CH₃)=CH₂), butenyl (C₄), pentenyl (C₅), and hexenyl (C₆).

10 C₂₋₁₂ alkynyl: The term "C₂₋₁₂ alkynyl" as used herein, pertains to an alkyl group having one or more carbon-carbon triple bonds.

Examples of unsaturated alkynyl groups include, but are not limited to, ethynyl (ethinyl, -C≡CH) and 2-propynyl (propargyl, -CH₂-C≡CH).

15 C₃₋₁₂ cycloalkyl: The term "C₃₋₁₂ cycloalkyl" as used herein, pertains to an alkyl group which is also a cyclyl group; that is, a monovalent moiety obtained by removing a hydrogen atom from an alicyclic ring atom of a cyclic hydrocarbon (carbocyclic) compound, which moiety has from 3 to 7 carbon atoms, including from 3 to 7
20 ring atoms.

Examples of cycloalkyl groups include, but are not limited to, those derived from:

saturated monocyclic hydrocarbon compounds:

25 cyclopropane (C₃), cyclobutane (C₄), cyclopentane (C₅), cyclohexane (C₆), cycloheptane (C₇), methylcyclopropane (C₄), dimethylcyclopropane (C₅), methylcyclobutane (C₅), dimethylcyclobutane (C₆), methylcyclopentane (C₆), dimethylcyclopentane (C₇) and methylcyclohexane (C₇);

30 unsaturated monocyclic hydrocarbon compounds:

cyclopropene (C₃), cyclobutene (C₄), cyclopentene (C₅), cyclohexene (C₆), methylcyclopropene (C₄), dimethylcyclopropene (C₅), methylcyclobutene (C₅), dimethylcyclobutene (C₆), methylcyclopentene (C₆), dimethylcyclopentene (C₇) and
35 methylcyclohexene (C₇); and

saturated polycyclic hydrocarbon compounds:

norcarane (C₇), norpinane (C₇), norbornane (C₇).

C₃₋₂₀ heterocyclyl: The term "C₃₋₂₀ heterocyclyl" as used herein, pertains to a monovalent moiety obtained by removing a hydrogen atom from a ring atom of a heterocyclic compound, which moiety has from 3 to 20 ring atoms, of which from 1 to 10 are ring heteroatoms. Preferably, each ring has from 3 to 7 ring atoms, of which from 1 to 4 are ring heteroatoms.

In this context, the prefixes (e.g. C₃₋₂₀, C₃₋₇, C₅₋₆, etc.) denote the number of ring atoms, or range of number of ring atoms, whether carbon atoms or heteroatoms. For example, the term "C₅₋₆heterocyclyl", as used herein, pertains to a heterocyclyl group having 5 or 6 ring atoms.

15

Examples of monocyclic heterocyclyl groups include, but are not limited to, those derived from:

- N₁: aziridine (C₃), azetidine (C₄), pyrrolidine (tetrahydropyrrole) (C₅), pyrroline (e.g., 3-pyrroline, 2,5-dihydropyrrole) (C₅), 2H-pyrrole or 3H-pyrrole (isopyrrole, isoazole) (C₅), piperidine (C₆), dihydropyridine (C₆), tetrahydropyridine (C₆), azepine (C₇);
- O₁: oxirane (C₃), oxetane (C₄), oxolane (tetrahydrofuran) (C₅), oxole (dihydrofuran) (C₅), oxane (tetrahydropyran) (C₆), dihydropyran (C₆), pyran (C₆), oxepin (C₇);
- S₁: thiirane (C₃), thietane (C₄), thiolane (tetrahydrothiophene) (C₅), thiane (tetrahydrothiopyran) (C₆), thiepane (C₇);
- O₂: dioxolane (C₅), dioxane (C₆), and dioxepane (C₇);
- O₃: trioxane (C₆);
- N₂: imidazolidine (C₅), pyrazolidine (diazolidine) (C₅), imidazoline (C₅), pyrazoline (dihydropyrazole) (C₅), piperazine (C₆);
- N₁O₁: tetrahydrooxazole (C₅), dihydrooxazole (C₅), tetrahydroisoxazole (C₅), dihydroisoxazole (C₅), morpholine (C₆), tetrahydrooxazine (C₆), dihydrooxazine (C₆), oxazine (C₆);
- N₁S₁: thiazoline (C₅), thiazolidine (C₅), thiomorpholine (C₆);
- N₂O₁: oxadiazine (C₆);
- O₁S₁: oxathiole (C₅) and oxathiane (thioxane) (C₆); and,

N₁O₁S₁: oxathiazine (C₆).

Examples of substituted monocyclic heterocyclyl groups include those derived from saccharides, in cyclic form, for example, furanoses (C₅), such as arabinofuranose, lyxofuranose, ribofuranose, and xylofuranse, and pyranoses (C₆), such as allopyranose, altropyranose, glucopyranose, mannopyranose, gulopyranose, idopyranose, galactopyranose, and talopyranose.

C₅₋₂₀ aryl: The term "C₅₋₂₀ aryl", as used herein, pertains to a monovalent moiety obtained by removing a hydrogen atom from an aromatic ring atom of an aromatic compound, which moiety has from 3 to 20 ring atoms. Preferably, each ring has from 5 to 7 ring atoms.

In this context, the prefixes (e.g. C₃₋₂₀, C₅₋₇, C₅₋₆, etc.) denote the number of ring atoms, or range of number of ring atoms, whether carbon atoms or heteroatoms. For example, the term "C₅₋₆ aryl" as used herein, pertains to an aryl group having 5 or 6 ring atoms.

The ring atoms may be all carbon atoms, as in "carboaryl groups". Examples of carboaryl groups include, but are not limited to, those derived from benzene (i.e. phenyl) (C₆), naphthalene (C₁₀), azulene (C₁₀), anthracene (C₁₄), phenanthrene (C₁₄), naphthacene (C₁₈), and pyrene (C₁₆).

Examples of aryl groups which comprise fused rings, at least one of which is an aromatic ring, include, but are not limited to, groups derived from indane (e.g. 2,3-dihydro-1H-indene) (C₉), indene (C₉), isoindene (C₉), tetraline (1,2,3,4-tetrahydronaphthalene (C₁₀), acenaphthene (C₁₂), fluorene (C₁₃), phenalene (C₁₃), acephenanthrene (C₁₅), and aceanthrene (C₁₆).

Alternatively, the ring atoms may include one or more heteroatoms, as in "heteroaryl groups". Examples of monocyclic heteroaryl groups include, but are not limited to, those derived from:

N₁: pyrrole (azole) (C₅), pyridine (azine) (C₆);
O₁: furan (oxole) (C₅);
S₁: thiophene (thiole) (C₅);
N₁O₁: oxazole (C₅), isoxazole (C₅), isoxazine (C₆);
5 N₂O₁: oxadiazole (furazan) (C₅);
N₃O₁: oxatriazole (C₅);
N₁S₁: thiazole (C₅), isothiazole (C₅);
N₂: imidazole (1,3-diazole) (C₅), pyrazole (1,2-diazole) (C₅),
pyridazine (1,2-diazine) (C₆), pyrimidine (1,3-diazine) (C₆) (e.g.,
10 cytosine, thymine, uracil), pyrazine (1,4-diazine) (C₆);
N₃: triazole (C₅), triazine (C₆); and,
N₄: tetrazole (C₅).

15 Examples of heteroaryl which comprise fused rings, include, but are not limited to:

C₉ (with 2 fused rings) derived from benzofuran (O₁),
isobenzofuran (O₁), indole (N₁), isoindole (N₁), indolizine (N₁),
indoline (N₁), isoindoline (N₁), purine (N₄) (e.g., adenine,
guanine), benzimidazole (N₂), indazole (N₂), benzoxazole (N₁O₁),
20 benzisoxazole (N₁O₁), benzodioxole (O₂), benzofurazan (N₂O₁),
benzotriazole (N₃), benzothiofuran (S₁), benzothiazole (N₁S₁),
benzothiadiazaole (N₂S);

C₁₀ (with 2 fused rings) derived from chromene (O₁),
isochromene (O₁), chroman (O₁), isochroman (O₁), benzodioxan (O₂),
25 quinoline (N₁), isoquinoline (N₁), quinolizine (N₁), benzoxazine
(N₁O₁), benzodiazine (N₂), pyridopyridine (N₂), quinoxaline (N₂),
quinazoline (N₂), cinnoline (N₂), phthalazine (N₂), naphthyridine
(N₂), pteridine (N₄);

C₁₁ (with 2 fused rings) derived from benzodiazepine (N₂);

30 C₁₃ (with 3 fused rings) derived from carbazole (N₁),
dibenzofuran (O₁), dibenzothiophene (S₁), carboline (N₂), perimidine
(N₂), pyridoindole (N₂); and,

C₁₄ (with 3 fused rings) derived from acridine (N₁), xanthene
(O₁), thioxanthene (S₁), oxanthrene (O₂), phenoxathiin (O₁S₁),
35 phenazine (N₂), phenoxazine (N₁O₁), phenothiazine (N₁S₁), thianthrene
(S₂), phenanthridine (N₁), phenanthroline (N₂), phenazine (N₂).

The above groups, whether alone or part of another substituent, may themselves optionally be substituted with one or more groups selected from themselves and the additional substituents listed below.

Halo: -F, -Cl, -Br, and -I.

Hydroxy: -OH.

Ether: -OR, wherein R is an ether substituent, for example, a C₁₋₇ alkyl group (also referred to as a C₁₋₇ alkoxy group, discussed below), a C₃₋₂₀ heterocyclyl group (also referred to as a C₃₋₂₀ heterocyclyloxy group), or a C₅₋₂₀ aryl group (also referred to as a C₅₋₂₀ aryloxy group), preferably a C₁₋₇alkyl group.

Alkoxy: -OR, wherein R is an alkyl group, for example, a C₁₋₇ alkyl group. Examples of C₁₋₇ alkoxy groups include, but are not limited to, -OMe (methoxy), -OEt (ethoxy), -O(nPr) (n-propoxy), -O(iPr) (isopropoxy), -O(nBu) (n-butoxy), -O(sBu) (sec-butoxy), -O(iBu) (isobutoxy), and -O(tBu) (tert-butoxy).

Acetal: -CH(OR¹)(OR²), wherein R¹ and R² are independently acetal substituents, for example, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a C₁₋₇ alkyl group, or, in the case of a "cyclic" acetal group, R¹ and R², taken together with the two oxygen atoms to which they are attached, and the carbon atoms to which they are attached, form a heterocyclic ring having from 4 to 8 ring atoms. Examples of acetal groups include, but are not limited to, -CH(OMe)₂, -CH(OEt)₂, and -CH(OMe)(OEt).

Hemiacetal: -CH(OH)(OR¹), wherein R¹ is a hemiacetal substituent, for example, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a C₁₋₇ alkyl group. Examples of hemiacetal groups include, but are not limited to, -CH(OH)(OMe) and -CH(OH)(OEt).

Ketal: $-\text{CR}(\text{OR}^1)(\text{OR}^2)$, where R^1 and R^2 are as defined for acetals, and R is a ketal substituent other than hydrogen, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples ketal groups include, but are not limited to, $-\text{C}(\text{Me})(\text{OMe})_2$, $-\text{C}(\text{Me})(\text{OEt})_2$, $-\text{C}(\text{Me})(\text{OMe})(\text{OEt})$, $-\text{C}(\text{Et})(\text{OMe})_2$, $-\text{C}(\text{Et})(\text{OEt})_2$, and $-\text{C}(\text{Et})(\text{OMe})(\text{OEt})$.

Hemiketal: $-\text{CR}(\text{OH})(\text{OR}^1)$, where R^1 is as defined for hemiacetals, and R is a hemiketal substituent other than hydrogen, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of hemiacetal groups include, but are not limited to, $-\text{C}(\text{Me})(\text{OH})(\text{OMe})$, $-\text{C}(\text{Et})(\text{OH})(\text{OMe})$, $-\text{C}(\text{Me})(\text{OH})(\text{OEt})$, and $-\text{C}(\text{Et})(\text{OH})(\text{OEt})$.

Oxo (keto, -one): $=\text{O}$.

Thione (thioketone): $=\text{S}$.

Imino (imine): $=\text{NR}$, wherein R is an imino substituent, for example, hydrogen, C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably hydrogen or a C_{1-7} alkyl group. Examples of ester groups include, but are not limited to, $=\text{NH}$, $=\text{NMe}$, $=\text{NEt}$, and $=\text{NPh}$.

Formyl (carbaldehyde, carboxaldehyde): $-\text{C}(=\text{O})\text{H}$.

Acyl (keto): $-\text{C}(=\text{O})\text{R}$, wherein R is an acyl substituent, for example, a C_{1-7} alkyl group (also referred to as C_{1-7} alkylacyl or C_{1-7} alkanoyl), a C_{3-20} heterocyclyl group (also referred to as C_{3-20} heterocyclylacyl), or a C_{5-20} aryl group (also referred to as C_{5-20} arylacyl), preferably a C_{1-7} alkyl group. Examples of acyl groups include, but are not limited to, $-\text{C}(=\text{O})\text{CH}_3$ (acetyl), $-\text{C}(=\text{O})\text{CH}_2\text{CH}_3$ (propionyl), $-\text{C}(=\text{O})\text{C}(\text{CH}_3)_3$ (t-butyryl), and $-\text{C}(=\text{O})\text{Ph}$ (benzoyl, phenone).

Carboxy (carboxylic acid): $-\text{C}(=\text{O})\text{OH}$.

Thiocarboxy (thiocarboxylic acid): $-\text{C}(=\text{S})\text{SH}$.

Thiolocarboxy (thiolocarboxylic acid): $-\text{C}(=\text{O})\text{SH}$.

Thionocarboxy (thionocarboxylic acid): $-\text{C}(=\text{S})\text{OH}$.

Imidic acid: $-\text{C}(=\text{NH})\text{OH}$.

Hydroxamic acid: $-\text{C}(=\text{NOH})\text{OH}$.

Ester (carboxylate, carboxylic acid ester, oxycarbonyl): $-\text{C}(=\text{O})\text{OR}$, wherein R is an ester substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of ester groups include, but are not limited to, $-\text{C}(=\text{O})\text{OCH}_3$, $-\text{C}(=\text{O})\text{OCH}_2\text{CH}_3$, $-\text{C}(=\text{O})\text{OC}(\text{CH}_3)_3$, and $-\text{C}(=\text{O})\text{OPh}$.

Acyloxy (reverse ester): $-\text{OC}(=\text{O})\text{R}$, wherein R is an acyloxy substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of acyloxy groups include, but are not limited to, $-\text{OC}(=\text{O})\text{CH}_3$ (acetoxy), $-\text{OC}(=\text{O})\text{CH}_2\text{CH}_3$, $-\text{OC}(=\text{O})\text{C}(\text{CH}_3)_3$, $-\text{OC}(=\text{O})\text{Ph}$, and $-\text{OC}(=\text{O})\text{CH}_2\text{Ph}$.

Oxycarboxyloxy: $-\text{OC}(=\text{O})\text{OR}$, wherein R is an ester substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of ester groups include, but are not limited to, $-\text{OC}(=\text{O})\text{OCH}_3$, $-\text{OC}(=\text{O})\text{OCH}_2\text{CH}_3$, $-\text{OC}(=\text{O})\text{OC}(\text{CH}_3)_3$, and $-\text{OC}(=\text{O})\text{OPh}$.

Amino: $-\text{NR}^1\text{R}^2$, wherein R^1 and R^2 are independently amino substituents, for example, hydrogen, a C_{1-7} alkyl group (also referred to as C_{1-7} alkylamino or di- C_{1-7} alkylamino), a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably H or a C_{1-7} alkyl group, or, in the case of a "cyclic" amino group, R^1 and R^2 , taken together with the nitrogen atom to which they are attached,


form a heterocyclic ring having from 4 to 8 ring atoms. Amino groups may be primary ($-\text{NH}_2$), secondary ($-\text{NHR}^1$), or tertiary ($-\text{NHR}^1\text{R}^2$), and in cationic form, may be quaternary ($-\text{NR}^1\text{R}^2\text{R}^3$).

5 Examples of amino groups include, but are not limited to, $-\text{NH}_2$, $-\text{NHCH}_3$, $-\text{NHC}(\text{CH}_3)_2$, $-\text{N}(\text{CH}_3)_2$, $-\text{N}(\text{CH}_2\text{CH}_3)_2$, and $-\text{NHPh}$. Examples of cyclic amino groups include, but are not limited to, aziridino, azetidino, pyrrolidino, piperidino, piperazino, morpholino, and thiomorpholino.

10 Amido (carbamoyl, carbamyl, aminocarbonyl, carboxamide): $-\text{C}(=\text{O})\text{NR}^1\text{R}^2$, wherein R^1 and R^2 are independently amino substituents, as defined for amino groups. Examples of amido groups include, but are not limited to, $-\text{C}(=\text{O})\text{NH}_2$, $-\text{C}(=\text{O})\text{NHCH}_3$, $-\text{C}(=\text{O})\text{N}(\text{CH}_3)_2$, $-\text{C}(=\text{O})\text{NHCH}_2\text{CH}_3$, and $-\text{C}(=\text{O})\text{N}(\text{CH}_2\text{CH}_3)_2$, as well as amido groups in
15 which R^1 and R^2 , together with the nitrogen atom to which they are attached, form a heterocyclic structure as in, for example, piperidinocarbonyl, morpholinocarbonyl, thiomorpholinocarbonyl, and piperazinocarbonyl.

20 Thioamido (thiocarbamyl): $-\text{C}(=\text{S})\text{NR}^1\text{R}^2$, wherein R^1 and R^2 are independently amino substituents, as defined for amino groups. Examples of amido groups include, but are not limited to, $-\text{C}(=\text{S})\text{NH}_2$, $-\text{C}(=\text{S})\text{NHCH}_3$, $-\text{C}(=\text{S})\text{N}(\text{CH}_3)_2$, and $-\text{C}(=\text{S})\text{NHCH}_2\text{CH}_3$.

25 Acylamido (acylamino): $-\text{NR}^1\text{C}(=\text{O})\text{R}^2$, wherein R^1 is an amide substituent, for example, hydrogen, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably hydrogen or a C_{1-7} alkyl group, and R^2 is an acyl substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group,
30 preferably hydrogen or a C_{1-7} alkyl group. Examples of acylamide groups include, but are not limited to, $-\text{NHC}(=\text{O})\text{CH}_3$, $-\text{NHC}(=\text{O})\text{CH}_2\text{CH}_3$, and $-\text{NHC}(=\text{O})\text{Ph}$. R^1 and R^2 may together form a cyclic structure, as in, for example, succinimidyl, maleimidyl, and phthalimidyl:




Aminocarbonyloxy: $-OC(=O)NR^1R^2$, wherein R^1 and R^2 are independently amino substituents, as defined for amino groups. Examples of aminocarbonyloxy groups include, but are not limited to, $-OC(=O)NH_2$, $-OC(=O)NHMe$, $-OC(=O)NMe_2$, and $-OC(=O)NEt_2$.

Ureido: $-N(R^1)CONR^2R^3$ wherein R^2 and R^3 are independently amino substituents, as defined for amino groups, and R^1 is a ureido substituent, for example, hydrogen, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably hydrogen or a C_{1-7} alkyl group. Examples of ureido groups include, but are not limited to, $-NHCONH_2$, $-NHCONHMe$, $-NHCONHEt$, $-NHCONMe_2$, $-NHCONEt_2$, $-NMeCONH_2$, $-NMeCONHMe$, $-NMeCONHEt$, $-NMeCONMe_2$, and $-NMeCONEt_2$.

Guanidino: $-NH-C(=NH)NH_2$.

Tetrazolyl: a five membered aromatic ring having four nitrogen atoms and one carbon atom,



Imino: $=NR$, wherein R is an imino substituent, for example, for example, hydrogen, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably H or a C_{1-7} alkyl group. Examples of imino groups include, but are not limited to, $=NH$, $=NMe$, and $=NEt$.

Amidine (amidino): $-C(=NR)NR_2$, wherein each R is an amidine substituent, for example, hydrogen, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably H or a C_{1-7}

alkyl group. Examples of amidine groups include, but are not limited to, $-C(=NH)NH_2$, $-C(=NH)NMe_2$, and $-C(=NMe)NMe_2$.

Nitro: $-NO_2$.

5

Nitroso: $-NO$.

Azido: $-N_3$.

10 Cyano (nitrile, carbonitrile): $-CN$.

Isocyano: $-NC$.

Cyanato: $-OCN$.

15

Isocyanato: $-NCO$.

Thiocyano (thiocyanato): $-SCN$.

20 Isothiocyano (isothiocyanato): $-NCS$.

Sulfhydryl (thiol, mercapto): $-SH$.

25 Thioether (sulfide): $-SR$, wherein R is a thioether substituent, for example, a C_{1-7} alkyl group (also referred to as a C_{1-7} alkylthio group), a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of C_{1-7} alkylthio groups include, but are not limited to, $-SCH_3$ and $-SCH_2CH_3$.

30 Disulfide: $-SS-R$, wherein R is a disulfide substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group (also referred to herein as C_{1-7} alkyl disulfide). Examples of C_{1-7} alkyl disulfide groups include, but are not limited to, $-SSCH_3$ and $-SSCH_2CH_3$.

35

5 Sulfine (sulfinyl, sulfoxide): $-S(=O)R$, wherein R is a sulfine substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfine groups include, but are not limited to, $-S(=O)CH_3$ and $-S(=O)CH_2CH_3$.

10 Sulfone (sulfonyl): $-S(=O)_2R$, wherein R is a sulfone substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group, including, for example, a fluorinated or perfluorinated C_{1-7} alkyl group. Examples of sulfone groups include, but are not limited to, $-S(=O)_2CH_3$ (methanesulfonyl, mesyl), $-S(=O)_2CF_3$ (triflyl), $-S(=O)_2CH_2CH_3$ (esyl), $-S(=O)_2C_4F_9$ (nonafllyl), $-S(=O)_2CH_2CF_3$ (tresyl), $-S(=O)_2CH_2CH_2NH_2$ (tauryl), $-S(=O)_2Ph$ (phenylsulfonyl, besyl), 4-methylphenylsulfonyl (tosyl), 15 4-chlorophenylsulfonyl (closyl), 4-bromophenylsulfonyl (brosyl), 4-nitrophenyl (nosyl), 2-naphthalenesulfonate (napsyl), and 5-dimethylamino-naphthalen-1-ylsulfonate (dansyl).

20 Sulfinic acid (sulfino): $-S(=O)OH$, $-SO_2H$.

Sulfonic acid (sulfo): $-S(=O)_2OH$, $-SO_3H$.

25 Sulfinate (sulfinic acid ester): $-S(=O)OR$; wherein R is a sulfinate substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfinate groups include, but are not limited to, $-S(=O)OCH_3$ (methoxysulfinyl; methyl sulfinate) and $-S(=O)OCH_2CH_3$ (ethoxysulfinyl; ethyl sulfinate).

30 Sulfonate (sulfonic acid ester): $-S(=O)_2OR$, wherein R is a sulfonate substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfonate groups include, but are not limited to, $-S(=O)_2OCH_3$ (methoxysulfonyl; methyl sulfonate) and 35 $-S(=O)_2OCH_2CH_3$ (ethoxysulfonyl; ethyl sulfonate).

Sulfinyloxy: $-\text{OS}(=\text{O})\text{R}$, wherein R is a sulfinyloxy substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfinyloxy groups include, but are not limited to, $-\text{OS}(=\text{O})\text{CH}_3$ and

5 $-\text{OS}(=\text{O})\text{CH}_2\text{CH}_3$.

Sulfonyloxy: $-\text{OS}(=\text{O})_2\text{R}$, wherein R is a sulfonyloxy substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfonyloxy groups include, but are not limited to, $-\text{OS}(=\text{O})_2\text{CH}_3$ (mesylate) and

10 $-\text{OS}(=\text{O})_2\text{CH}_2\text{CH}_3$ (esylate).

Sulfate: $-\text{OS}(=\text{O})_2\text{OR}$; wherein R is a sulfate substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably a C_{1-7} alkyl group. Examples of sulfate groups include, but are not limited to, $-\text{OS}(=\text{O})_2\text{OCH}_3$ and $-\text{SO}(=\text{O})_2\text{OCH}_2\text{CH}_3$.

15

Sulfamyl (sulfamoyl; sulfinic acid amide; sulfinamide): $-\text{S}(=\text{O})\text{NR}^1\text{R}^2$, wherein R^1 and R^2 are independently amino substituents, as defined for amino groups. Examples of sulfamyl groups include, but are not limited to, $-\text{S}(=\text{O})\text{NH}_2$, $-\text{S}(=\text{O})\text{NH}(\text{CH}_3)$, $-\text{S}(=\text{O})\text{N}(\text{CH}_3)_2$, $-\text{S}(=\text{O})\text{NH}(\text{CH}_2\text{CH}_3)$, $-\text{S}(=\text{O})\text{N}(\text{CH}_2\text{CH}_3)_2$, and $-\text{S}(=\text{O})\text{NHPh}$.

20

Sulfonamido (sulfinamoyl; sulfonic acid amide; sulfonamide): $-\text{S}(=\text{O})_2\text{NR}^1\text{R}^2$, wherein R^1 and R^2 are independently amino substituents, as defined for amino groups. Examples of sulfonamido groups include, but are not limited to, $-\text{S}(=\text{O})_2\text{NH}_2$, $-\text{S}(=\text{O})_2\text{NH}(\text{CH}_3)$, $-\text{S}(=\text{O})_2\text{N}(\text{CH}_3)_2$, $-\text{S}(=\text{O})_2\text{NH}(\text{CH}_2\text{CH}_3)$, $-\text{S}(=\text{O})_2\text{N}(\text{CH}_2\text{CH}_3)_2$, and $-\text{S}(=\text{O})_2\text{NHPh}$.

25

Sulfamino: $-\text{NR}^1\text{S}(=\text{O})_2\text{OH}$, wherein R^1 is an amino substituent, as defined for amino groups. Examples of sulfamino groups include, but are not limited to, $-\text{NHS}(=\text{O})_2\text{OH}$ and $-\text{N}(\text{CH}_3)\text{S}(=\text{O})_2\text{OH}$.

30

Sulfonamino: $-\text{NR}^1\text{S}(=\text{O})_2\text{R}$, wherein R^1 is an amino substituent, as defined for amino groups, and R is a sulfonamino substituent, for example, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl

35

group, preferably a C₁₋₇ alkyl group. Examples of sulfonamino groups include, but are not limited to, -NHS(=O)₂CH₃ and -N(CH₃)S(=O)₂C₆H₅.

- 5 Sulfinamino: -NR¹S(=O)R, wherein R¹ is an amino substituent, as defined for amino groups, and R is a sulfinamino substituent, for example, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a C₁₋₇ alkyl group. Examples of sulfinamino groups include, but are not limited to, -NHS(=O)CH₃ and
10 -N(CH₃)S(=O)C₆H₅.

- Phosphino (phosphine): -PR₂, wherein R is a phosphino substituent, for example, -H, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably -H, a C₁₋₇ alkyl group, or a C₅₋₂₀ aryl
15 group. Examples of phosphino groups include, but are not limited to, -PH₂, -P(CH₃)₂, -P(CH₂CH₃)₂, -P(t-Bu)₂, and -P(Ph)₂.

Phospho: -P(=O)₂.

- 20 Phosphinyl (phosphine oxide): -P(=O)R₂, wherein R is a phosphinyl substituent, for example, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably a C₁₋₇ alkyl group or a C₅₋₂₀ aryl group. Examples of phosphinyl groups include, but are not limited to, -P(=O)(CH₃)₂, -P(=O)(CH₂CH₃)₂, -P(=O)(t-Bu)₂, and
25 -P(=O)(Ph)₂.

Phosphonic acid (phosphono): -P(=O)(OH)₂.

- 30 Phosphonate (phosphono ester): -P(=O)(OR)₂, where R is a phosphonate substituent, for example, -H, a C₁₋₇ alkyl group, a C₃₋₂₀ heterocyclyl group, or a C₅₋₂₀ aryl group, preferably -H, a C₁₋₇ alkyl group, or a C₅₋₂₀ aryl group. Examples of phosphonate groups include, but are not limited to, -P(=O)(OCH₃)₂, -P(=O)(OCH₂CH₃)₂, -P(=O)(O-t-Bu)₂, and -P(=O)(OPh)₂.

35

Phosphoric acid (phosphonooxy): -OP(=O)(OH)₂.

Phosphate (phosphonooxy ester): $-\text{OP}(=\text{O})(\text{OR})_2$, where R is a phosphate substituent, for example, -H, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably -H, a C_{1-7} alkyl group, or a C_{5-20} aryl group. Examples of phosphate groups include, but are not limited to, $-\text{OP}(=\text{O})(\text{OCH}_3)_2$, $-\text{OP}(=\text{O})(\text{OCH}_2\text{CH}_3)_2$, $-\text{OP}(=\text{O})(\text{O}-t\text{-Bu})_2$, and $-\text{OP}(=\text{O})(\text{OPh})_2$.

Phosphorous acid: $-\text{OP}(\text{OH})_2$.

10

Phosphite: $-\text{OP}(\text{OR})_2$, where R is a phosphite substituent, for example, -H, a C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably -H, a C_{1-7} alkyl group, or a C_{5-20} aryl group. Examples of phosphite groups include, but are not limited to, $-\text{OP}(\text{OCH}_3)_2$, $-\text{OP}(\text{OCH}_2\text{CH}_3)_2$, $-\text{OP}(\text{O}-t\text{-Bu})_2$, and $-\text{OP}(\text{OPh})_2$.

15

Phosphoramidite: $-\text{OP}(\text{OR}^1)-\text{NR}^2_2$, where R^1 and R^2 are phosphoramidite substituents, for example, -H, a (optionally substituted) C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably -H, a C_{1-7} alkyl group, or a C_{5-20} aryl group. Examples of phosphoramidite groups include, but are not limited to, $-\text{OP}(\text{OCH}_2\text{CH}_3)-\text{N}(\text{CH}_3)_2$, $-\text{OP}(\text{OCH}_2\text{CH}_3)-\text{N}(i\text{-Pr})_2$, and $-\text{OP}(\text{OCH}_2\text{CH}_2\text{CN})-\text{N}(i\text{-Pr})_2$.

20

Phosphoramidate: $-\text{OP}(=\text{O})(\text{OR}^1)-\text{NR}^2_2$, where R^1 and R^2 are phosphoramidate substituents, for example, -H, a (optionally substituted) C_{1-7} alkyl group, a C_{3-20} heterocyclyl group, or a C_{5-20} aryl group, preferably -H, a C_{1-7} alkyl group, or a C_{5-20} aryl group. Examples of phosphoramidate groups include, but are not limited to, $-\text{OP}(=\text{O})(\text{OCH}_2\text{CH}_3)-\text{N}(\text{CH}_3)_2$, $-\text{OP}(=\text{O})(\text{OCH}_2\text{CH}_3)-\text{N}(i\text{-Pr})_2$, and $-\text{OP}(=\text{O})(\text{OCH}_2\text{CH}_2\text{CN})-\text{N}(i\text{-Pr})_2$.

25

30

Alkylene

C_{3-12} alkylene: The term " C_{3-12} alkylene", as used herein, pertains to a bidentate moiety obtained by removing two hydrogen atoms, either both from the same carbon atom, or one from each of two

35

different carbon atoms, of a hydrocarbon compound having from 3 to 12 carbon atoms (unless otherwise specified), which may be aliphatic or alicyclic, and which may be saturated, partially unsaturated, or fully unsaturated. Thus, the term "alkylene" includes the sub-classes alkenylene, alkynylene, cycloalkylene, etc., discussed below.

Examples of linear saturated C₃₋₁₂ alkylene groups include, but are not limited to, -(CH₂)_n- where n is an integer from 3 to 12, for example, -CH₂CH₂CH₂- (propylene), -CH₂CH₂CH₂CH₂- (butylene), -CH₂CH₂CH₂CH₂CH₂- (pentylene) and -CH₂CH₂CH₂CH₂CH₂CH₂CH₂- (heptylene).

Examples of branched saturated C₃₋₁₂ alkylene groups include, but are not limited to, -CH(CH₃)CH₂-, -CH(CH₃)CH₂CH₂-, -CH(CH₃)CH₂CH₂CH₂-, -CH₂CH(CH₃)CH₂-, -CH₂CH(CH₃)CH₂CH₂-, -CH(CH₂CH₃)-, -CH(CH₂CH₃)CH₂-, and -CH₂CH(CH₂CH₃)CH₂-.

Examples of linear partially unsaturated C₃₋₁₂ alkylene groups (C₃₋₁₂ alkenylene, and alkynylene groups) include, but are not limited to, -CH=CH-CH₂-, -CH₂-CH=CH₂-, -CH=CH-CH₂-CH₂-, -CH=CH-CH₂-CH₂-CH₂-, -CH=CH-CH=CH-, -CH=CH-CH=CH-CH₂-, -CH=CH-CH=CH-CH₂-CH₂-, -CH=CH-CH₂-CH=CH-, -CH=CH-CH₂-CH₂-CH=CH-, and -CH₂-C≡C-CH₂-.

Examples of branched partially unsaturated C₃₋₁₂ alkylene groups (C₃₋₁₂ alkenylene and alkynylene groups) include, but are not limited to, -C(CH₃)=CH-, -C(CH₃)=CH-CH₂-, -CH=CH-CH(CH₃)- and -C≡C-CH(CH₃)-.

Examples of alicyclic saturated C₃₋₁₂ alkylene groups (C₃₋₁₂ cycloalkylenes) include, but are not limited to, cyclopentylene (e.g. cyclopent-1,3-ylene), and cyclohexylene (e.g. cyclohex-1,4-ylene).

Examples of alicyclic partially unsaturated C₃₋₁₂ alkylene groups (C₃₋₁₂ cycloalkylenes) include, but are not limited to, cyclopentenylene (e.g. 4-cyclopenten-1,3-ylene), cyclohexenylene

(e.g. 2-cyclohexen-1,4-ylene; 3-cyclohexen-1,2-ylene; 2,5-cyclohexadien-1,4-ylene).

Proliferative Diseases

5 One of ordinary skill in the art is readily able to determine whether or not a candidate compound treats a proliferative condition for any particular cell type. For example, assays which may conveniently be used to assess the activity offered by a particular compound are described in the examples below.

10

The term "proliferative disease" pertains to an unwanted or uncontrolled cellular proliferation of excessive or abnormal cells which is undesired, such as, neoplastic or hyperplastic growth, whether *in vitro* or *in vivo*.

15

Examples of proliferative conditions include, but are not limited to, benign, pre-malignant, and malignant cellular proliferation, including but not limited to, neoplasms and tumours (e.g. histocytoma, glioma, astrocyoma, osteoma), cancers (e.g. lung cancer, small cell lung cancer, gastrointestinal cancer, bowel cancer, colon cancer, breast carcinoma, ovarian carcinoma, prostate cancer, testicular cancer, liver cancer, kidney cancer, bladder cancer, pancreas cancer, brain cancer, sarcoma, osteosarcoma, Kaposi's sarcoma, melanoma), leukemias, psoriasis, bone diseases, fibroproliferative disorders (e.g. of connective tissues), and atherosclerosis.

20

25

Any type of cell may be treated, including but not limited to, lung, gastrointestinal (including, e.g. bowel, colon), breast (mammary), ovarian, prostate, liver (hepatic), kidney (renal), bladder, pancreas, brain, and skin.

30

Methods of Treatment

As described above, the present invention provide the use of a compound of formula III in a method of therapy. Also provided is a method of treatment, comprising administering to a subject in need

35

of treatment a therapeutically-effective amount of a compound of formula III, preferably in the form of a pharmaceutical composition, which is the third aspect of the present invention. The term "therapeutically effective amount" is an amount sufficient to show benefit to a patient. Such benefit may be at least amelioration of at least one symptom. The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated. Prescription of treatment, e.g. decisions on dosage, is within the responsibility of general practitioners and other medical doctors.

A compound may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated. Examples of treatments and therapies include, but are not limited to, chemotherapy (the administration of active agents, including, e.g. drugs; surgery; and radiation therapy. If the compound of formula III bears a carbamate-based nitrogen protecting group which may be removed *in vivo*, then the methods of treatment described in WO 00/12507 (ADEPT, GDEPT and PDT) may be used.

Pharmaceutical compositions according to the present invention, and for use in accordance with the present invention, may comprise, in addition to the active ingredient, i.e. a compound of formula III, a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material will depend on the route of administration, which may be oral, or by injection, e.g. cutaneous, subcutaneous, or intravenous.

Pharmaceutical compositions for oral administration may be in tablet, capsule, powder or liquid form. A tablet may comprise a solid carrier or an adjuvant. Liquid pharmaceutical compositions generally comprise a liquid carrier such as water, petroleum,

animal or vegetable oils, mineral oil or synthetic oil.
Physiological saline solution, dextrose or other saccharide
solution or glycols such as ethylene glycol, propylene glycol or
polyethylene glycol may be included. A capsule may comprise a
5 solid carrier such as gelatin.

For intravenous, cutaneous or subcutaneous injection, or injection
at the site of affliction, the active ingredient will be in the
form of a parenterally acceptable aqueous solution which is
10 pyrogen-free and has suitable pH, isotonicity and stability. Those
of relevant skill in the art are well able to prepare suitable
solutions using, for example, isotonic vehicles such as Sodium
Chloride Injection, Ringer's Injection, Lactated Ringer's
Injection. Preservatives, stabilisers, buffers, antioxidants
15 and/or other additives may be included, as required.

Includes Other Forms


Unless otherwise specified, included in the above are the well
known ionic, salt, solvate, and protected forms of these
20 substituents. For example, a reference to carboxylic acid ($-\text{COOH}$)
also includes the anionic (carboxylate) form ($-\text{COO}^-$), a salt or
solvate thereof, as well as conventional protected forms.
Similarly, a reference to an amino group includes the protonated
form ($-\text{N}^+\text{HR}^1\text{R}^2$), a salt or solvate of the amino group, for example,
25 a hydrochloride salt, as well as conventional protected forms of an
amino group. Similarly, a reference to a hydroxyl group also
includes the anionic form ($-\text{O}^-$), a salt or solvate thereof, as well
as conventional protected forms.

Isomers, Salts and Solvates

Certain compounds may exist in one or more particular geometric,
optical, enantiomeric, diasteriomeric, epimeric, atropic,
stereoisomeric, tautomeric, conformational, or anomeric forms,
including but not limited to, cis- and trans-forms; E- and Z-forms;
35 c-, t-, and r- forms; endo- and exo-forms; R-, S-, and meso-forms;
D- and L-forms; d- and l-forms; (+) and (-) forms; keto-, enol-,


and enolate-forms; syn- and anti-forms; synclinal- and anticlinal-forms; α - and β -forms; axial and equatorial forms; boat-, chair-, twist-, envelope-, and halfchair-forms; and combinations thereof, hereinafter collectively referred to as "isomers" (or "isomeric forms").

Preferably compounds of the present invention have the following stereochemistry at the C11 position:



Note that, except as discussed below for tautomeric forms, specifically excluded from the term "isomers", as used herein, are structural (or constitutional) isomers (i.e. isomers which differ in the connections between atoms rather than merely by the position of atoms in space). For example, a reference to a methoxy group, $-OCH_3$, is not to be construed as a reference to its structural isomer, a hydroxymethyl group, $-CH_2OH$. Similarly, a reference to ortho-chlorophenyl is not to be construed as a reference to its structural isomer, meta-chlorophenyl. However, a reference to a class of structures may well include structurally isomeric forms falling within that class (e.g. C_{1-7} alkyl includes n-propyl and iso-propyl; butyl includes n-, iso-, sec-, and tert-butyl; methoxyphenyl includes ortho-, meta-, and para-methoxyphenyl).

The above exclusion does not pertain to tautomeric forms, for example, keto-, enol-, and enolate-forms, as in, for example, the following tautomeric pairs: keto/enol (illustrated below), imine/enamine, amide/imino alcohol, amidine/amidine, nitroso/oxime, thioketone/enethiol, N-nitroso/hydroxyazo, and nitro/aci-nitro.



Note that specifically included in the term "isomer" are compounds with one or more isotopic substitutions. For example, H may be in any isotopic form, including ^1H , ^2H (D), and ^3H (T); C may be in any isotopic form, including ^{12}C , ^{13}C , and ^{14}C ; O may be in any isotopic form, including ^{16}O and ^{18}O ; and the like.

Unless otherwise specified, a reference to a particular compound includes all such isomeric forms, including (wholly or partially) racemic and other mixtures thereof. Methods for the preparation (e.g. asymmetric synthesis) and separation (e.g. fractional crystallisation and chromatographic means) of such isomeric forms are either known in the art or are readily obtained by adapting the methods taught herein, or known methods, in a known manner.

Unless otherwise specified, a reference to a particular compound also includes ionic, salt, solvate, and protected forms of thereof, for example, as discussed below.

It may be convenient or desirable to prepare, purify, and/or handle a corresponding salt of the active compound, for example, a pharmaceutically-acceptable salt. Examples of pharmaceutically acceptable salts are discussed in Berge, et al., *J. Pharm. Sci.*, **66**, 1-19 (1977).

For example, if the compound is anionic, or has a functional group which may be anionic (e.g. $-\text{COOH}$ may be $-\text{COO}^-$), then a salt may be formed with a suitable cation. Examples of suitable inorganic cations include, but are not limited to, alkali metal ions such as Na^+ and K^+ , alkaline earth cations such as Ca^{2+} and Mg^{2+} , and other cations such as Al^{3+} . Examples of suitable organic cations include, but are not limited to, ammonium ion (i.e. NH_4^+) and substituted ammonium ions (e.g. NH_3R^+ , NH_2R_2^+ , NHR_3^+ , NR_4^+). Examples of some suitable substituted ammonium ions are those derived from: ethylamine, diethylamine, dicyclohexylamine, triethylamine, butylamine, ethylenediamine, ethanolamine, diethanolamine,


piperazine, benzylamine, phenylbenzylamine, choline, meglumine, and tromethamine, as well as amino acids, such as lysine and arginine. An example of a common quaternary ammonium ion is $N(CH_3)_4^+$.

- 5 If the compound is cationic, or has a functional group which may be cationic (e.g. $-NH_2$ may be $-NH_3^+$), then a salt may be formed with a suitable anion. Examples of suitable inorganic anions include, but are not limited to, those derived from the following inorganic acids: hydrochloric, hydrobromic, hydroiodic, sulfuric, sulfurous, 10 nitric, nitrous, phosphoric, and phosphorous.

Examples of suitable organic anions include, but are not limited to, those derived from the following organic acids:

- 2-acetoxybenzoic, acetic, ascorbic, aspartic, benzoic, 15 camphorsulfonic, cinnamic, citric, edetic, ethanedisulfonic, ethanesulfonic, fumaric, glucheptonic, gluconic, glutamic, glycolic, hydroxymaleic, hydroxynaphthalene carboxylic, isethionic, lactic, lactobionic, lauric, maleic, malic, methanesulfonic, mucic, oleic, oxalic, palmitic, pantoic, pantothenic, phenylacetic, 20 phenylsulfonic, propionic, pyruvic, salicylic, stearic, succinic, sulfanilic, tartaric, toluenesulfonic, and valeric. Examples of suitable polymeric organic anions include, but are not limited to, those derived from the following polymeric acids: tannic acid, carboxymethyl cellulose.

25 A particular salt form of interest can be formed from compounds of formula III, where R^{10} and R^{16} form an imine bond, by reacting said compound with a bisulphite salt to form a bisulphite derivative of the PBD. These compounds can be represented as:




30 where M is a monovalent pharmaceutically acceptable cation, or if the compound is a dimer, the two M groups may together represent a

divalent pharmaceutically acceptable cation, and the other groups are as previously defined.

It may be convenient or desirable to prepare, purify, and/or handle a corresponding solvate of the active compound. The term "solvate" is used herein in the conventional sense to refer to a complex of solute (e.g. active compound, salt of active compound) and solvent. If the solvent is water, the solvate may be conveniently referred to as a hydrate, for example, a mono-hydrate, a di-hydrate, a tri-hydrate, etc.

Solvates of particular relevance to the present invention are those where the solvent adds across the imine bond of the PBD moiety, which is illustrated below where the solvent is water or an alcohol ($R^A OH$, where R^A is an ether substituent as described above):



These forms can be called the carbinolamine and carbinolamine ether forms of the PBD. The balance of these equilibria depend on the conditions in which the compounds are found, as well as the nature of the moiety itself.


In general any nucleophilic solvent is capable of forming such solvates as illustrated above for hydroxylic solvents. Other nucleophilic solvents include thiols and amines.

These solvates may be isolated in solid form, for example, by lyophilisation.

General synthetic routes

The synthesis of PBD compounds is extensively discussed in WO 00/12508, which discussion is incorporated herein by reference.

As discussed in that patent application, a key step in a preferred route to PBDs is a cyclisation to produce the B-ring, involving generation of an aldehyde (or functional equivalent thereof) at what will be the 11-position, and attack thereon by the Pro-N10-nitrogen:




IV

wherein the substituents are as defined in the second aspect of the invention. The "masked aldehyde" -CPQ may be an acetal or thioacetal, in which case the cyclisation involves unmasking. Alternatively, it may be an alcohol -CHOH, in which case the reaction involves oxidation, e.g. by means of TPAP, TEMPO or DMSO (Swern oxidation).


In this reaction, R^{12} is preferably $O-R^{14}$, i.e. a protected hydroxy group, and R^{13} is H.

The masked aldehyde compound can be produced by condensing a corresponding 2,4-substituted pyrrolidine with a 2-nitrobenzoic acid:



The nitro group can then be reduced to $-NH_2$ and protected by reaction with a suitable agent, e.g. a chloroformate, which provides the removable nitrogen protecting group in the compound of formula IV.

A process involving the oxidation-cyclization procedure is illustrated in scheme 1 (an alternative type of cyclisation will be described later with reference to scheme 2).



Scheme 1

Exposure of the alcohol (B) (in which the Pro-N10-nitrogen is generally protected as carbamate) to tetrapropylammonium perruthenate (TPAP)/N-methylmorpholine N-oxide (NMO) over A4 sieves results in oxidation accompanied by spontaneous B-ring closure to afford the desired product IV. The TPAP/NMO oxidation procedure is found to be particularly convenient for small scale reactions while the use of DMSO-based oxidation methods, particularly Swern oxidation, proves superior for larger scale work (e.g. > 1 g). A particularly preferred oxidising agent is (diacetoxyiodo)benzene (1.1 eq) and TEMPO (0.1 eq) dissolved in CH₂Cl₂.

The uncyclized alcohol (B) may be prepared by the reaction of a nitrogen protection reagent of formula D, which is preferably a

chloroformate or acid chloride, to a solution of the amino alcohol C, generally in solution, generally in the presence of a base such as pyridine (preferably 2 equivalents) at a moderate temperature (e.g. at 0°C). Under these conditions little or no O-acylation is usually observed.


The key amino alcohol C may be prepared by reduction of the corresponding nitro compound E, by choosing a method which will leave the rest of the molecule intact. Treatment of E with tin (II) chloride in a suitable solvent, e.g. refluxing methanol, generally affords, after the removal of the tin salts, the desired product in high yield.

Exposure of E to hydrazine/Raney nickel avoids the production of tin salts and may result in a higher yield of C, although this method is less compatible with the range of possible C and A-ring substituents. For instance, if there is C-ring unsaturation (either in the ring itself, or in R₂ or R₃), this technique may be unsuitable. Another suitable means of reduction would be catalytic hydrogenation using palladium on carbon as a catalyst.

The nitro compound of formula E may be prepared by coupling the appropriate o-nitrobenzoyl chloride to a compound of formula F, e.g. in the presence of K₂CO₃ at -25°C under a N₂ atmosphere. Compounds of formula F can be readily prepared, for example by olefination of the ketone derived from L-trans-hydroxy proline. The ketone intermediate can also be exploited by conversion to the enol triflate for use in palladium mediated coupling reactions.

The o-nitrobenzoyl chloride is synthesised from the o-nitrobenzoic acid (or alkyl ester after hydrolysis) of formula G, which itself is prepared from the vanillic acid (or alkyl ester) derivative H. Many of these are commercially available and some are disclosed in Althuis, T.H. and Hess, H.J., *J. Medicinal Chem.*, 20(1), 146-266 (1977).

Alternative Cyclisation (Scheme 2)




Scheme 2

- 5 In scheme 1, the final or penultimate step was an oxidative cyclisation. An alternative, using thioacetal coupling, is shown in scheme 2. Mercury-mediated unmasking causes cyclisation to the protected PBD compound IV.
- 10 The thioacetal compound may be prepared as shown in scheme 2: the thioacetal protected C-ring [prepared via a literature method: Langley, D.R. & Thurston, D.E., *J. Organic Chemistry*, 52, 91-97 (1987)] is coupled to the o-nitrobenzoic acid (or alkyl ester after hydrolysis) (G) using a literature procedure. The resulting nitro
- 15 compound cannot be reduced by hydrogenation, because of the thioacetal group, so the tin(II) chloride method is used to afford

the amine. This is then N-protected, e.g., by reaction with a chloroformate or acid chloride, such as 2,2,2-trichloroethylchloroformate.

- 5 Acetal-containing C-rings can be used as an alternative in this type of route with deprotection involving other methods, including the use of acidic conditions.

Dimer Synthesis (Scheme 3)



G'

Scheme 3

PBD dimers may be synthesized using the strategy developed for the synthesis of the protected PBD monomers. The synthesis routes illustrated in scheme 3 show compounds when the dimer linkage is of the formula -O-(CH₂)_n-O-. The step of dimer formation is normally

carried out to form a bis(nitro acid) G'. This compound can then be treated as compound G in either scheme 1 or scheme 2 above.

5 The bis(nitro acid) G' may be obtained by nitrating (e.g. using 70% nitric acid) the bis(carboxylic acid). This can be synthesised by alkylation of two equivalents of the relevant benzoic acid with the appropriate diiodoalkane under basic conditions. Many benzoic acids are commercially available and others can be synthesised by conventional methods. Alternatively, the relevant benzoic acid
10 esters can be joined together by a Mitsunobu etherification with an appropriate alkanediol, followed by nitration, and then hydrolysis (not illustrated).


15 An alternative synthesis of the bis(nitro acid) involves oxidation of the bis(nitro aldehyde), e.g. with potassium permanganate. This can be obtained in turn by direct nitration of the bis(aldehyde), e.g. with 70% HNO₃. Finally, the bis(aldehyde) can be obtained via the Mitsunobu etherification of two equivalents of the benzoic aldehyde with the appropriate alkanediol.

20

Alternative routes to PBDs

Alternative methods of synthesising N10 protected PBDs are disclosed in co-pending application PCT/GB2004/003873 (filed 10 September 2004) which claims priority from GB0321295.8 (filed 11
25 September 2003), which describes the use of isocyanate intermediates.

Formation of Compound of formula I



Following cyclisation to form the B-ring, the C11-alcohol IV is then preferably re-protected, by conventional means to provide IIb.

For example, if R^{11} is TBS, the protection can take place by reacting IV with TBSOTf and 2,6-lutidine. Cleavage of the C2-protecting group from IIb then provides the C2 alcohol. For example, where the C2 protecting group (R^{14}) is acyl, this deprotection may be performed by addition of an aqueous solution of K_2CO_3 .

This reprotection at the C11 position and deprotection of the C2 alcohol allows subsequent reaction of selectively the C2 alcohol position leaving the C11 position unaffected.


The C2-alcohol may then be oxidized to the ketone IIa. Preferably this oxidation is performed under Swern conditions, in good yield. However, other oxidation methods involving TPAP or the Dess Martin reagent also provide the ketone in good yield.

If R^2 in the compound of formula I is $-OSO_2CH_3$, $-OSO_2(C_nF_{2n+1})$ where $n=0,1$ or 4 , or $-OSO_2R^5$, then the conversion from IIb may be achieved by treatment with the appropriate anhydride. For example, if R^2 is triflate that reaction with trifluoromethanesulfonic anhydride in DCM in the presence of pyridine.

If R^2 in the compound of formula I is $-I$ or $-Br$, then the conversion from **IIb** may be achieved by reaction with hydrazine and iodine or bromine respectively.

- 5 If R^2 in the compound of formula I is $-Cl$, then the conversion from **IIb** may be achieved by reaction with a phosphorous oxychloride (e.g. $POCl_3$).

Synthesis of compounds of formula III



10

This compound of formula I may be reacted under a variety of conditions to yield PBD precursor molecules with pendant groups coupled at the C2 position **IIIc**.

- 15 In particular, the use of palladium catalysed coupling is preferred, such as Suzuki, Stille and Heck coupling. The palladium catalyst may be any suitable catalyst, for example $Pd(PPh_3)_4$, $Pd(OCOCH_3)_2$, $PdCl_2$, $Pd(dba)_3$. The compounds which are coupled may be any suitable reactant, e.g. for Heck, alkenes with an sp^2 H; for
- 20 Stille, organostannanes; and for Suzuki, organoboron derivatives.

In a preferred aspect of the invention, the coupling may be performed under microwave conditions. Typically, the palladium catalyst, such as $Pd(PPh_3)_4$, is solid supported, for example on

25 polystyrene, to facilitate work-up and allow potential recycling of

catalyst. Unreacted boronic acid can be sequestered following complete consumption of triflate using PS-DEAM, with a phase separator cartridge being used to isolate the coupling product. Such a method allows for the parallel synthesis of more than one
5 (e.g. up to 10, 20 or 30) compound at the same time.

The imine bond in the compound of formula IIIc can be unprotected by standard methods to yield the unprotected compound IIIa (which may be in its carbinolamine or carbinoilamine ether form ,
10 depending on the solvents used). For example if R¹⁰ is Alloc, then the deprotection is carried using palladium to remove the N10 protecting group, followed by the elimination of water. If R¹⁰ is Troc, then the deprotection is carried out using a Cd/Pb couple to yield the compound of formula IIIa.

15 If the nitrogen protecting group (R¹⁰) is such that the desired end product still contains it, e.g. if it is removable *in vivo*, then the compound of formula IIb may be synthesised by removal of the oxygen protecting groups under suitable conditions.

20 Further Preferences

The following preferences may apply to all aspects of the invention as described above, or may relate to a single aspect. The preferences may be combined together in any combination.

25 R⁶ to R⁹

If the compound is a dimer, it is preferred that the dimer bridge is of formula -O-(CH₂)_n-O-, where n is from 3 to 12, and more preferably 3 to 7. It is preferred that the substituents R⁸ join
30 to form the dimer bridge.

R⁹ is preferably H.

35 R⁶ is preferably selected from H, OH, OR, SH, NH₂, nitro and halo, and is more preferably H or halo, and most preferably is H.

R⁷ and R⁸ (when the compound is not a dimer) are preferably independently selected from H, OH, OR, SH, SR, NH₂, NHR, NRR', and halo, and more preferably independently selected from H, OH and OR, where R is preferably selected from optionally substituted C₁₋₇ alkyl, C₃₋₁₀ heterocyclyl and C₅₋₁₀ aryl groups. Particularly preferred substituents at the 7- and 8- positions are OMe and OCH₂Ph.

In the first aspect of the invention, R¹⁰ is preferably Troc. R¹¹ is preferably a silyl oxygen protecting group (more preferably TBS) or THP. R² is preferably triflate. Accordingly, in a particularly preferred embodiment of the first aspect of the invention, R¹⁰ is Troc, R¹¹ is TBS or THP and R² is triflate.

In the second aspect of the invention, R¹⁴ is preferably a methyl ester. R¹¹ is preferably a silyl oxygen protecting group (more preferably TBS) or THP. Accordingly, in a particularly preferred embodiment of the second aspect of the invention R¹⁴ is a methyl ester and R¹¹ is TBS or THP. Furthermore, R¹⁰ is preferably Troc.

In some embodiments of the third aspect of the invention, R¹⁰ is preferably Troc and R¹⁶ is O-R¹¹, wherein R¹¹ is preferably a silyl oxygen protecting group (more preferably TBS) or THP.

In other embodiments of the third aspect of the invention, R¹⁰ and R¹⁶ together form a double bond between N10 and C11.

In the third aspect of the invention R¹⁵ is preferably selected from optionally substituted C₅₋₂₀ aryl groups and optionally substituted C₁₋₇ alkyl groups, which group has a carbon-carbon double or triple bond conjugated to the double bond in the C-ring.

Novel compounds of the present invention preferably have R¹⁰ and R¹⁶ forming a double bond between N10 and C11. Preferably, the novel compounds of the invention are dimers through C8, i.e. the R⁸ groups of each monomer form together a dimer bridge having the

formula $-X-R''-X-$ linking the monomers. More preferably, the dimer bridge is of formula $-O-(CH_2)_n-O-$, where n is 3 to 12, more preferably 3, 5, or 7. The preferences for R^6 , R^7 and R^9 are as expressed above. R^{15} is preferably selected from:

- 5 (i) optionally substituted C_{5-20} aryl groups;
- (ii) substituted C_2 alkyl groups; and
- (iii) optionally substituted C_{3-7} alkyl groups.

10 In particular groups (ii) and (iii) above, preferably have a carbon-carbon double or triple bond conjugated to that between C_2 and C_3 .

Group (i) above are more preferably optionally substituted C_{5-7} aryl groups, and most preferably optionally substituted phenyl groups.

15 Group (ii) is preferably either a vinyl group substituted with an amido group, and more preferably with an amido group which is $-C(=O)N(CH_3)_2$; or a ethynyl group substituted with an optionally substituted C_{5-7} aryl group, more preferably phenyl.


20 Group (iii) is preferably an optionally substituted propylene group, for example $-CH=CH-CH_3$.

25 If R is optionally substituted C_{1-12} alkyl, it is preferred that it is optionally substituted C_{1-7} alkyl.

Examples**Example 1 - Formation of key dimer intermediate (2-**


[[[(trifluoromethyl)sulfonyl]oxy]-1,10,11,11a-tetrahydro-5H-
pyrrolo[2,1-c][1,4]benzodiazepin-5-one]] (13))

(2S,4R)-N-(Benzyloxycarbonyl)-2-t-butyldimethylsilyloxymethyl-4-
hydroxypyrrolidine (1)



Compound 1 is formed in high yield in a four step process known in
the art starting from *trans*-4-hydroxy-L-proline (S.J. Gregson et
al., *J. Med. Chem.*, 2004, 1161-1174).

(2S,4R)-N-(Benzyloxycarbonyl)-2-t-butyldimethylsilyloxymethyl-4-
oxyacetylpyrrolidine (2)




(a) pyridine, Ac₂O, DMAP, THF, 16 h, 96%;

Pyridine (18.3 g, 18.7 mL, 232 mmol, 1.1 eq), acetic anhydride
(23.6 g, 21.8 mL, 232 mmol, 1.1 eq) and DMAP (5.14 g, 42.1 mmol,
0.2 eq) were added to a stirred solution of 1 (76.9 g, 211 mmol) in
dry THF (1 L). The reaction mixture was stirred for 16 hours after
which time TLC (95:5 v/v CHCl₃/MeOH) showed the complete
consumption of starting material. Excess solvent was removed by
rotary evaporation and the residue was dissolved in EtOAc (1 L),
washed with 1N HCl (2 x 1 L), H₂O (1 L), brine (1 L) and dried
(MgSO₄). Filtration and evaporation of the solvent afforded
acetate 2 as a colourless oil (80.7 g, 94%): ¹H NMR (400 MHz, CDCl₃)
(rotamers) δ 7.36-7.12 (m, 5H), 5.30-5.10 (m, 3H), 4.09-3.97 (m,

2H), 3.74-3.55 (m, 3H), 2.36-2.29 (m, 1H), 2.11-2.06 (m, 1H), 2.02 (s, 3H), 0.87 (s, 6H), 0.86 (s, 3H), 0.03 and 0.00 (s x 2, 6H); MS (ES), m/z (relative intensity) 430 ($[M + Na]^+$, 95), 408 ($[M + H]^+$, 100).


5

(2*S*,4*R*)-2-*t*-Butyldimethylsilyloxymethyl-4-oxyacetylpyrrolidine (3)



A slurry of silyl ether 2 (1.95 g, 4.80 mmol) and 10% Pd/C (0.17 g) in absolute ethanol (10 mL) was subjected to Parr hydrogenation at 45 Psi for 16 h after which time TLC (95:5 v/v CHCl₃/MeOH) showed the complete consumption of starting material. The reaction mixture was filtered through celite to remove the Pd/C, and the filter pad was washed repeatedly with ethanol. Excess solvent was removed by rotary evaporation under reduced pressure to afford the amine 3 as a pale orange waxy oil (1.28 g, 98%): IR (CHCl₃) 3315, 2930, 2858, 1739, 1652, 1472, 1435, 1375, 1251, 1088, 838, 779, 667 cm⁻¹.

1,1'-[[(Propane-1,3-diyl)dioxy]bis[(2*S*,4*R*)-(5-methoxy-2-nitro-1,4-phenylene)carbonyl]]bis[2-(*tert*-butyldimethylsilyloxymethyl)-4-oxyacetylpyrrolidine] (5)




A catalytic amount of DMF (2 drops) was added to a stirred solution of the nitro-acid 4 (8.12 g, 17.4 mmol)¹ and oxalyl chloride (3.80 mL, 5.52 g, 43.5 mmol, 2.5 eq) in dry THF (250 mL). The initial precipitate dissolved gradually and the reaction mixture was allowed to stir for 16 h at room temperature. The resulting acid chloride solution was added dropwise to a stirred mixture of the amine 3 (11.9 g, 43.5 mmol, 2.5 eq), TEA (9.71 mL, 7.05 g, 69.7 mmol, 4.0 eq) and H₂O (2.26 mL) in THF (100 mL) at 0°C

(ice/acetone) under a nitrogen atmosphere. The reaction mixture was allowed to warm to room temperature and stirred for a further 2.5 h. Excess THF was removed by rotary evaporation and the resulting residue was partitioned between H₂O (400 mL) and EtOAc (400 mL). The layers were allowed to separate and the aqueous layer was extracted with EtOAc (3 x 200 mL). The combined organic layers were then washed with saturated NH₄Cl (200 mL), saturated NaHCO₃ (200 mL), brine (200 mL) and dried (MgSO₄). Filtration and evaporation of the solvent gave the crude product as a dark oil.

Purification by flash chromatography (99.7:0.3 v/v CHCl₃/MeOH) isolated the pure amide 5 as a light yellow glass (13.3 g, 78%): ¹H NMR (400 MHz, CDCl₃) δ 7.60 (s, 2H), 6.60 (s, 2H), 5.06 (br s, 2H), 4.44 (br s, 2H), 4.25-4.20 (m, 4H), 4.10-4.08 (m, 2H), 3.80 (s, 6H), 3.64-3.62 (m, 2H), 3.36-3.32 (m, 2H), 3.11-3.08 (m, 2H), 2.36-2.26 (m, 4H), 2.13-2.08 (m, 2H), 1.92 (s, 6H), 0.80 (s, 18H), 0.00 (s x 2, 12H); ¹³C NMR (100.6 MHz, CDCl₃) δ 171.0, 166.3, 154.5, 148.2, 137.4, 128.0, 127.2, 109.2, 108.5, 72.9, 65.6, 62.6, 57.4, 56.5, 54.8, 33.0, 28.6, 25.8, 21.0, 18.1; MS (ES), m/z (relative intensity) 1000 ([M + Na]⁺, 39), 978 ([M + H]⁺, 63), 977 (M⁺, 100), 812 (13).


1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(2S,4R)-(5-methoxy-2-amino-1,4-phenylene)carbonyl]]bis[2-(tert-butyldimethylsilyloxymethyl)-4-oxyacetylpyrrolidine] (6)



Sodium dithionite (16.59 g, 95.27 mmol, 5 eq) was added to a stirred solution of amide 5 (18.6 g, 19.1 mmol) in H₂O (200 mL) and THF (400 mL). The reaction mixture was allowed to stir for 36 h after which time excess THF was removed by rotary evaporation and the resulting residue was extracted with EtOAc (3 x 250 mL). The combined organic layers were then washed with H₂O (300 mL), brine (300 mL) and dried (MgSO₄). Filtration and evaporation of the

solvent yielded the crude product which was purified by flash chromatography (80:20 v/v hexane/EtOAc then gradient to neat EtOAc) to afford the product 6 as a yellow foam (9.53 g, 55%): ^1H NMR (400 MHz, CDCl_3) (rotamers) δ 6.70 and 6.67 (s x 2, 2H), 6.25 and 6.23 (s x 2, 2H), 5.20 (br s, 2H), 4.49 (br s, 4H), 4.16-4.05 (m, 6H), 3.70 (s, 6H), 3.68-3.57 (m, 4H), 2.36-2.27 (m, 4H), 2.12-2.04 (m, 2H), 1.96 (s, 6H), 0.85 (s, 18H), 0.01 and 0.00 (s x 2, 12H); ^{13}C NMR (100.6 MHz, CDCl_3) δ 170.6, 170.0, 141.1, 116.3, 113.1, 102.3, 102.1, 102.0, 66.2, 65.3, 65.2, 57.0, 28.9, 18.2; MS (ES), m/z (relative intensity) 946 ($\text{M}^+ + 29$, 43), 933 ($[\text{M} + 16]^+$, 61), 932 ($[\text{M} + 15]^+$, 100), 918 ($[\text{M} + \text{H}]^+$, 72).


1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(2*S*,4*R*)-[5-methoxy-1,4-phenylene-2-(2,2,2-trichloroethoxycarbonylamino)]carbonyl]]bis[2-(tert-butyldimethylsilyloxymethyl)-4-oxyacetylpyrrolidine]] (7)



A solution of 2,2,2-trichloroethyl chloroformate (3.58 mL, 5.50 g, 26.0 mmol, 2.2 eq) in dry DCM (60 mL) was added dropwise to a solution of anhydrous pyridine (3.82 mL, 3.80 g, 47.2 mmol, 4.0 eq) and bis-aniline 6 (10.8 g, 11.8 mmol) in dry DCM (150 mL) at -10°C (liq. N_2 /ethanediol). After 16 h at room temperature, the reaction mixture was washed with saturated NH_4Cl (2 x 150 mL), saturated CuSO_4 (150 mL), H_2O (150 mL), brine (150 mL) and dried (MgSO_4). Filtration and evaporation of the solvent yielded a yellow viscous oil which was purified by flash chromatography (70:30 v/v hexane/EtOAc) to afford the product 7 as a white glass (13.8 g, 92%): ^1H NMR (400 MHz, CDCl_3) δ 9.42 (br s, 1H), 7.83 (s, 2H), 6.76 and 6.74 (s x 2, 2H), 5.21 (br s, 2H), 4.79 and 4.73 (d x 2, 4H, $J = 12.0$ Hz), 4.56 (br s, 2H), 4.26-4.23 (m, 4H), 4.09-4.04 (m, 2H), 3.74 (s, 6H), 3.72-3.68 (m, 2H), 3.60 (br s, 4H), 2.40-2.32 (m, 4H), 2.23-2.08 (m, 2H), 1.95 (s, 6H), 0.85 (s, 18H), 0.01 and 0.00

(s x 2, 12H); ^{13}C NMR (100.6 MHz, CDCl_3) δ 170.4, 169.2, 151.9, 151.5, 150.8, 143.4, 132.6, 114.4, 111.7, 95.3, 74.4, 65.5, 65.4, 57.3, 56.4, 32.5, 28.8, 25.8, 21.1, 18.1, 14.9; MS (ES), m/z (relative intensity) 1306 ($[\text{M} + 38]^+$, 92), 1304 ($[\text{M} + 36]^+$, 100),
 5 1282 ($[\text{M} + 14]^+$, 97), 1280 ($[\text{M} + 12]^+$, 55) .

1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(2*S*,4*R*)-[5-methoxy-1,4-phenylene-2-(2,2,2-trichloroethoxycarbonylamino)]carbonyl]]bis(2-hydroxymethyl-4-oxyacetylpyrrolidine) (8)




10

A mixture of glacial acetic acid (310 mL) and H_2O (100 mL) was added to a solution of 7 (13.8 g, 10.9 mmol) in THF (250 mL) and was stirred for 16 h at room temperature. The reaction mixture was diluted with DCM (750 mL) and neutralised with saturated NaHCO_3 (5
 15 L). The aqueous layer was extracted with DCM (3 x 500 mL) and the organic layers were combined, washed with brine (1 L) and dried (MgSO_4). TLC (60:40 v/v hexane/EtOAc) revealed the complete disappearance of the starting material. Filtration and evaporation of the solvent afforded the crude product which was purified by
 20 flash column chromatography (99.7:0.3 v/v $\text{CHCl}_3/\text{MeOH}$ then gradient to 96:4 v/v $\text{CHCl}_3/\text{MeOH}$) to provide the product 8 as a white glass (11.6 g, > 100%): ^1H NMR (500 MHz, CDCl_3) δ 8.92 (br s, 2H), 7.55 (s, 1H), 6.71 (s, 1H), 5.18 (br s, 2H), 4.78 (d, 2H, $J = 12.0$ Hz), 4.72 (d, 2H, $J = 12.0$ Hz), 4.50 (br s, 2H), 4.22-4.19 (m, 4H), 4.00 (br s, 2H), 3.78 (s, 6H), 3.76-3.52 (m, 6H), , 2.32-2.30 (m, 2H), 2.21-2.17 (m, 2H), 2.09-2.04 (m, 2H) 1.94 (s, 6H); ^{13}C NMR (125.8
 25 MHz, CDCl_3) δ 170.4, 152.2, 149.8, 145.0, 111.3, 106.5, 95.6, 74.4, 72.5, 65.4, 64.1, 58.7, 56.5, 56.3, 33.6, 29.1, 21.1.


30

1,1'-[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS,2R) -10-(2,2,2-trichloroethoxycarbonyl)-11-hydroxy-7-methoxy-2-oxyacetyl-1,2,3,10,11,11a-hexahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]] (9)




TEMPO (0.69 g, 4.42 mmol, 0.4 eq) and BAIB (15.7 g, 48.7 mmol, 4.4 eq) were added to a stirred solution of diol 8 (11.5 g, 11.1 mmol) in DCM (150 mL). The reaction mixture was allowed to stir for 2 h and diluted with DCM (400 mL), washed with saturated NaHSO₃ (500 mL), saturated NaHCO₃ (500 mL), brine (200 mL) and dried (MgSO₄). Filtration and evaporation of the solvent afforded the crude product which was purified by flash column chromatography (99.9:0.1 v/v CHCl₃/MeOH then gradient to 99.7:0.3 v/v CHCl₃/MeOH) to provide the product 9 as a light yellow glass (4.43 g, 39%): ¹H NMR (400 MHz, CDCl₃) δ 7.28 (s, 2H, H6), 6.84 (s, 2H, H9), 5.68 (d, 2H, J = 9.1 Hz, H11), 5.37-5.35 (m, 2H, H2), 5.18 (d, 2H, J = 12.0 Hz, Troc CH₂), 4.32-4.21 (m, 6H, OCH₂CH₂CH₂O, Troc CH₂), 4.03 (dd, 2H, J = 13.2, 2.6 Hz, H3), 3.92 (s, 6H, OCH₃ x 2), 3.39-3.69 (m, 4H, H3 and H11a), 2.39-2.35 (m, 6H, OCH₂CH₂CH₂O and H1), 2.03 (s, 6H, CH₃CO₂ x 2); ¹³C NMR (100.6 MHz, CDCl₃) δ 170.4 (CH₃CO₂), 167.4 (C_{quat}), 154.3 (C_{quat}), 150.5 (C_{quat}), 149.1 (C_{quat}), 127.4 (C_{quat}), 124.9 (C_{quat}), 114.1 (C9), 110.9 (C6), 95.0 (Troc CCl₃), 87.5 (C11), 75.0 (Troc CH₂), 71.4 (C2), 65.5 (OCH₂CH₂CH₂O), 58.4 (C11a), 56.1 (OCH₃), 51.1 (C3), 35.8 (C1), 29.1 (OCH₂CH₂CH₂O), 21.0 (CH₃CO₂); MS (ES), m/z (relative intensity) 1058 ([M + Na]⁺, 100).

1,1'-[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS,2R)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butyldimethylsilyloxy)-7-methoxy-2-oxyacetyl-1,2,3,10,11,11a-hexahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one]] (10)




TBSOTf (2.70 mL, 3.10 g, 11.7 mmol, 3.0 eq) was added to a stirred solution of bis-alcohol 9 (4.05 g, 3.91 mmol) and 2,6-lutidine (1.82 mL, 1.68 g, 15.6 mmol, 4.0 eq) in DCM (50 mL). The reaction mixture was allowed to stir for 2.5 h and diluted with DCM (150 mL), washed with saturated CuSO₄ (2 X 100 mL), saturated NaHCO₃ (100 mL), brine (200 mL) and dried (MgSO₄). Filtration and evaporation of the solvent afforded the crude product which was purified by flash column chromatography (99.9:0.1 v/v CHCl₃/MeOH) to provide the product 10 as a white glass (5.05 g, >100%): ¹H NMR (400 MHz, CDCl₃) δ 7.05 (s, 2H, H₆), 6.52 (s, 2H, H₉), 5.53 (d, 2H, J = 9.0 Hz, H₁₁), 5.14 (br s, 2H, H₂), 4.99 (d, 2H, J = 12.0 Hz, Troc CH₂), 4.06-3.87 (m, 8H, OCH₂CH₂CH₂O, Troc CH₂ and H_{11a}), 3.71 (s, 6H, OCH₃ x 2), 3.48-3.43 (m, 4H, H₃), 2.21-2.11 (m, 4H, OCH₂CH₂CH₂O and H₁), 2.03-1.96 (m, 2H, H₁), 1.81 (s, 6H, CH₃CO₂ x 2), 0.63 (s, 18H, TBS CH₃ x 6), 0.00 (s x 2, 12H, TBS CH₃ x 4); ¹³C NMR (100.6 MHz, CDCl₃) δ 170.3 (CH₃CO₂), 167.9 (C_{quat}), 153.6 (C_{quat}), 150.4 (C_{quat}), 149.2 (C_{quat}), 127.9 (C_{quat}), 125.5 (C_{quat}), 113.9 (C₉), 110.7 (C₆), 95.2 (Troc CCl₃), 88.2 (C₁₁), 74.7 (Troc CH₂), 71.7 (C₂), 65.0 (OCH₂CH₂CH₂O), 60.5 (C_{11a}), 56.1 (OCH₃), 51.2 (C₃), 36.2 (C₁), 28.8 (OCH₂CH₂CH₂O), 25.6 (TBS CH₃), 21.0 (CH₃CO₂), 17.8 (TBS C_{quat}), 14.2 and 14.1 (TBS CH₃); MS (ES), m/z (relative intensity) 1285 ([M + 21]⁺, 100), 1265 ([M + H]⁺, 75).

1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11*S*,11*aS*,2*R*)-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-hydroxy-1,2,3,10,11,11*a*-hexahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]] (11)



A solution of K_2CO_3 (93 mg, 0.67 mmol, 5.0 eq) in H_2O (2 mL) was added dropwise to a stirred solution of acetate 10 (170 mg, 0.13 mmol) in MeOH (3 mL). The initial colorless solution eventually turned yellow and the formation of a white precipitate was observed. The reaction mixture was allowed to stir for 16 h when TLC (95:5 v/v $CHCl_3$ /MeOH) showed the complete consumption of the starting material. Excess solvent was removed by rotary evaporation and the mixture was carefully neutralized with 1N HCl to pH 7. The resulting mixture was extracted with EtOAc (3 x 25 mL) and the combined organic layers were then washed with brine (40 mL) and dried ($MgSO_4$). Filtration and removal of the solvent afforded the product 11 as a white glass (151 mg, 95%): 1H NMR (400 MHz, $CDCl_3$) δ 6.94 (s, 2H, H6), 6.52 (s, 2H, H9), 5.53 (d, 2H, J = 9.0 Hz, H11), 5.00 (d, 2H, J = 12.0 Hz, Troc CH_2), 4.36-4.35 (m, 2H, H2), 4.06-3.82 (m, 8H, $OCH_2CH_2CH_2O$, Troc CH_2 and H3), 3.61 (s, 6H, OCH_3 x 2), 3.54-3.48 (m, 2H, H11a), 3.39-3.34 (m, 2H, H3), 2.96 and 2.95 (br s x 2, 2H, OH x 2), 2.21-2.20 (m, 2H, $OCH_2CH_2CH_2O$), 2.19-2.08 (m, 2H, H1), 1.90-1.74 (m, 2H, H1), 0.64 (s, 18H, TBS CH_3 x 6), 0.00 (s, 12H, TBS CH_3 x 4); ^{13}C NMR (100.6 MHz, $CDCl_3$) δ 168.5 (C_{quat}), 153.6 (C_{quat}), 150.3 (C_{quat}), 149.1 (C_{quat}), 127.9 (C_{quat}), 125.4 (C_{quat}), 113.9 (C9), 110.7 (C6), 95.2 (Troc CCl_3), 88.3 (C11), 74.7 (Troc CH_2), 69.4 (C2), 65.0 ($OCH_2CH_2CH_2O$), 60.9 (C11a), 55.9 (OCH_3), 54.1 (C3), 38.8 (C1), 28.9 ($OCH_2CH_2CH_2O$), 25.6 (TBS CH_3), 17.8 (TBS C_{quat}); MS (ES), m/z (relative intensity) 1196 ($[M + 16]^+$, 100), 1181 ($[M + H]^+$, 82).


1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11*S*,11*aS*)-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-oxo-1,2,3,10,11,11*a*-hexahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]] (12)



A solution of anhydrous DMSO (0.82 mL, 0.90 g, 11.5 mmol, 6.0 eq) in dry DCM (20 mL) was added dropwise to a stirred solution of oxalyl chloride (2.88 mL of a 2 M solution in DCM, 5.76 mmol, 3.0 eq) under a nitrogen atmosphere at -60°C (liq N_2/CHCl_3). After stirring at -55°C for 1.5 h, a solution of the substrate 11 (2.26 g, 1.92 mmol) in dry DCM (30 mL) was added dropwise to the reaction mixture, which was then stirred for a further 2 h at -45°C . A solution of TEA (10.8 mL, 7.82 g; 71.7 mmol, 4.2 eq) in dry DCM (90 mL) was added dropwise to the mixture and stirred for a further 30 min. The reaction mixture was left to warm to 0°C , washed with cold 1 N HCl (2 x 50 mL), H_2O (50 mL), brine (50 mL) and dried (MgSO_4). Filtration and evaporation of the solvent *in vacuo* afforded the crude product which was purified by flash column chromatography (70:30 v/v hexane/EtOAc then gradient to 40:60 v/v hexane/EtOAc) to afford carbinolamine 12 as a white glass (1.62 g, 72%): ^1H NMR (400 MHz, CDCl_3) δ 7.02 (s, 2H, H6), 6.54 (s, 2H, H9), 5.59 (d, 2H, $J = 9.2$ Hz, H11), 4.98 (d, 2H, $J = 12.0$ Hz, Troc CH_2), 4.09–3.86 (m, 8H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$, Troc CH_2 and H3), 3.75–3.66 (m, 10H, OCH_3 x 2, H11a, and H3), 2.72 (dd, 2H, $J = 10.2, 19.6$ Hz, H1), 2.82 (dd, 2H, $J = 2.6, 19.6$ Hz, H1), 2.22–2.19 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 0.63 (s, 18H, TBS CH_3 x 6), 0.00 (s x 2, 12H, TBS CH_3 x 4); ^{13}C NMR (100.6 MHz, CDCl_3) δ 207.7 (C2), 168.0 (C_{quat}), 153.7 (C_{quat}), 150.7 (C_{quat}), 149.4 (C_{quat}), 127.8 (C_{quat}), 124.6 (C_{quat}), 114.0 (C9), 110.6 (C6), 95.1 (Troc CCl_3), 87.4 (C11), 74.8 (Troc CH_2), 65.0 ($\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 58.9 (C11a), 56.1 (OCH_3), 53.0 (C3), 40.3 (C1), 28.8 ($\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 25.6 (TBS CH_3), 17.8 (TBS C_{quat}); MS (ES), m/z

(relative intensity) 1224 ($[M + 48]^+$, 100), 1210 ($[M + 34]^+$, 60), 1199 ($[M + Na]^+$, 35), 1192 ($[M + 16]^+$, 40), 1176 (M^+ , 18).

1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11*S*,11*aS*)-10-(2,2,2-trichloroethoxycarbonyl)-11-(*tert*-butyldimethylsilyloxy)-7-methoxy-2-[[[(trifluoromethyl)sulfonyl]oxy]-1,10,11,11*a*-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one]] (13)




Anhydrous triflic anhydride (3.09 mL, 5.19 g, 18.4 mmol, 22 eq) taken from a freshly opened ampule was added rapidly in one portion to a vigorously stirred solution of ketone 12 (0.98 g, 0.84 mmol) and anhydrous pyridine (1.49 mL, 1.46 g, 18.4 mmol, 22 eq) in dry DCM (50 mL) at room temperature under a nitrogen atmosphere. The initial precipitate dissolved gradually and the solution eventually turned a dark red colour. The reaction mixture was allowed to stir for 4.5 h when TLC (80:20 v/v EtOAc/hexane) revealed the complete consumption of the starting material. The mixture was poured into cold saturated NaHCO_3 (60 mL) and extracted with DCM (3 x 80 mL). The combined organic layers were then washed with saturated CuSO_4 (2 x 125 mL), brine (125 mL) and dried (MgSO_4). Filtration and evaporation of the solvent afforded the crude product which was purified by flash column chromatography (80:20 v/v hexane/EtOAc) to afford triflate 13 as a light yellow glass (0.74 mg, 61%): $[\alpha]_D^{25} = +46.0^\circ$ ($c = 0.33$, CHCl_3); $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.23 (s, 2H, H6), 7.19 (s, 2H, H3), 6.77 (s, 2H, H9), 5.94 (d, 2H, $J = 8.9$ Hz, H11), 5.23 (d, 2H, $J = 12.0$ Hz, Troc CH_2), 4.31-4.28 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 4.18 (d, 2H, $J = 12.2$ Hz, Troc CH_2), 4.15-4.13 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 3.95-3.91 (m, 8H, $\text{OCH}_3 \times 2$, H11a), 3.35 (dd, 2H, $J = 11.0, 16.6$ Hz, H1), 2.84 (d, 2H, $J = 16.6$ Hz, H1), 2.46-2.44 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 0.89 (s, 18H, TBS $\text{CH}_3 \times 6$), 0.29 and 0.26 (s x 2, 12H, TBS $\text{CH}_3 \times 4$); $^{13}\text{C NMR}$ (100.6 MHz, CDCl_3) δ 164.9 (C_{quat}), 153.6 (C_{quat}), 151.0 (C_{quat}), 149.5 (C_{quat}), 136.0 (C_{quat}), 127.7 (C_{quat}), 123.9

(C_{quat}), 121.0 (C3), 114.0 (C9), 110.9 (C6), 95.1 (Troc CCl₃), 86.3 (C11), 74.8 (Troc CH₂), 65.0 (OCH₂CH₂CH₂O), 60.6 (C11a), 56.2 (OCH₃), 34.4 (C1), 28.8 (OCH₂CH₂CH₂O), 25.6 (TBS CH₃), 17.8 (TBS C_{quat}); IR (CHCl₃) 3020, 2957, 2860, 1725, 1674, 1651, 1604, 1516, 1466, 1454, 1431, 1409, 1329, 1312, 1274, 1216, 1138, 1113, 1083, 1042, 1006, 900, 840, 757, 668, 646, 610 cm⁻¹; MS (ES), m/z (relative intensity) 1461 ([M + 21]⁺, 100), 1440 (M⁺, 55).

Example 2 - 1,1'-[[(Propane-1,3-diyl)dioxy]bis[(11aS)-7-methoxy-2-[(N,N-dimethylaminocarbonyl)vinyl]-1,11a-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]] (ZC-204)


1,1'-[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butyldimethylsilyloxy)-7-methoxy-2-[(N,N-dimethylaminocarbonyl)vinyl]-1,10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]] (14)



A mixture of triflate 13 (732 mg, 1.22 mmol), *N,N*-dimethylacrylamide (0.14 mL, 134 mg, 1.35 mmol, 5.0 eq), DABCO (152 mg, 1.35 mmol, 5.0 eq), (CH₃CN)₂PdCl₂ (14 mg, 0.05 mmol, 0.2 eq) and MeOH (15 mL) was stirred at 55 – 65 °C for 16 h. The reaction was worked-up by pouring the mixture into CHCl₃ (20 mL) and aqueous NaHCO₃ (20 mL). The aqueous layer was extracted with CHCl₃ (3 x 20 mL) and the CHCl₃ extracts were combined, washed with H₂O (50 mL), brine (50 mL) and dried (MgSO₄). Filtration and evaporation of solvent gave the crude product as a dark brown glass. The residue was purified by flash column chromatography (99.9:0.1 v/v CHCl₃/MeOH then gradient to 99.2:0.8 v/v CHCl₃/MeOH) to give the coupled product 14 as a light yellow glass (80 mg, 22%): [α]_D²⁶ = +87.2 ° (c = 0.11, CHCl₃); ¹H NMR (250 MHz, CDCl₃) δ 7.51 (d, 2H, J = 15.0 Hz, H12), 7.32 (s, 2H, H3), 7.28 (s, 2H, H6), 6.79 (s, 2H, H9), 6.10 (d, 2H, J = 15.0 Hz, H13), 5.88 (d, 2H, J = 8.8 Hz, H11),

5.23 (d, 2H, $J = 12.0$ Hz, Troc CH_2), 4.33-4.28 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$),
 4.18 (d, 1H, $J = 12.2$ Hz, Troc CH_2), 4.18-4.14 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$),
 3.98-3.95 (m, 8H, H11a and $\text{OCH}_3 \times 2$), 3.17-3.06 (m, 8H, H1 and NCH_3
 $\times 2$), 2.65 (d, 1H, $J = 16.2$ Hz, H1), 2.47-2.44 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$),
 0.92 (s, 18H, TBS $\text{CH}_3 \times 6$), 0.29 and 0.27 (s $\times 2$, 12H, TBS $\text{CH}_3 \times 4$);
 ^{13}C NMR (62.9 MHz, CDCl_3) δ 166.5 (CONMe₂), 132.5 (C12), 132.6 (C3),
 116.8 (C13), 114.0 (C9), 110.8 (C6), 95.2 (Troc CCl_3), 86.9 (C11),
 74.8 (Troc CH_2), 65.0 ($\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 61.9 (C11a), 56.2 (OCH_3), 33.5
 (C1), 28.8 ($\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 25.6 (TBS CH_3), 17.9 (TBS C_{quat}); MS (ES),
 m/z (relative intensity) 1337 ($[\text{M} - \text{H}]^+$, 100), 1327 (27), 1316
 (34).

1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11aS)-7-methoxy-2-[(*N,N*-
 dimethylaminocarbonyl)vinyl]-1,11a-dihydro-5H-pyrrolo[2,1-
 c][1,4]benzodiazepine-5-one]]] (ZC-204)




10% Cd/Pd couple (120 mg, 0.99 mmol, 16.5 eq) was added to a
 rapidly stirring mixture of 14 (79 mg, 0.06 mmol), THF (1.5 mL) and
 1 N NH_4OAc (1.5 mL). The reaction mixture was allowed to stir for
 3.5 h. The solids were filtered and rinsed with H_2O and CHCl_3 . The
 aqueous layer was extracted with CHCl_3 (3 \times 20 mL), and the organic
 extracts were combined, washed with brine (50 mL) and dried
 (MgSO_4). Filtration and evaporation of solvent left a yellow solid
 which was purified by flash column chromatography (99.9:0.1 v/v
 $\text{CHCl}_3/\text{MeOH}$ then gradient to 95:5 v/v $\text{CHCl}_3/\text{MeOH}$) to afford ZC-204 as
 a yellow glass (13.4 mg, 31%): $[\alpha]^{25}_D = +235.7^\circ$ ($c = 0.07$, CHCl_3);
 ^1H NMR (400 MHz, CDCl_3) δ 7.88 (d, 2H, $J = 3.9$ Hz, H11), 7.54 (d,
 2H, $J = 15.1$ Hz, H12), 7.51 (s, 2H, H3), 7.34 (s, 2H, H6), 6.88 (s,
 2H, H9), 6.18 (d, 2H, $J = 15.0$ Hz, H13), 4.43-4.27 (m, 6H,
 $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$ and H11a), 3.95 (s, 6H, $\text{OCH}_3 \times 2$), 3.41-3.34 (m, 2H,
 H1), 3.23-3.04 (m, 14H, H1 and $\text{NCH}_3 \times 4$), 2.47-2.44 (m, 2H,
 $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$); ^{13}C NMR (62.9 MHz, CDCl_3) δ 166.4 (CONMe₂), 162.1 (C11),

161.7 (C_{quat}), 151.5 (C_{quat}), 148.2 (C_{quat}), 140.4 (C_{quat}), 135.2 (C12),
 132.2 (C3), 121.8 (C_{quat}), 118.6 (C_{quat}), 117.0 (C13), 112.0 (C9),
 111.3 (C6), 65.4 (OCH₂CH₂CH₂O), 56.2 (OCH₃), 54.2 (C11a), 37.4 and
 35.9 (NCH₃), 33.8 (C1) 28.8 (OCH₂CH₂CH₂O); MS (ES), m/z (relative
 intensity) 741 ([M + H₂O]⁺, 25), 723 (M⁺, 62).

Example 3 - 1,1'-[[(Propane-1,3-diyl)dioxy]bis[(11aS)-7-methoxy-2-(p-methoxybenzene)-1,11a-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]] (ZC-207)


1,1'-[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butyldimethylsilyloxy)-7-methoxy-2-(p-methoxybenzene)-1,10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]] (15)



A solution of TEA (0.20 mL, 148 mg, 1.46 mmol, 6.0 eq) in H₂O (1.5 mL) and EtOH (10 mL) was added to a solution of triflate 13 (350 mg, 0.24 mmol) in toluene (10 mL) at room temperature. To this mixture 4-methoxybenzeneboronic acid (96 mg, 0.63 mmol, 2.6 eq) and Pd(PPh₃)₄ (11 mg, 9 μmol, 0.04 eq) were added. The reaction mixture was allowed to stir for 15 min when TLC (80:20 v/v EtOAc/hexane) revealed the complete consumption of the starting material. Excess solvent was removed and the residue was dissolved in EtOAc (25 mL), washed with H₂O (15 mL), brine (15 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (80:20 v/v hexane/EtOAc then gradient to 50:50 v/v hexane/EtOAc) to afford 15 as a yellow glass (286 mg, 87%): ¹H NMR (400 MHz, CDCl₃) δ 7.38 (s, 2H, H3), 7.32-7.28 (m, 6H, H6 and H13), 6.92 (d, 4H, J = 8.7 Hz, H14), 6.81 (s, 2H, H9), 5.93 (d, 2H, J = 8.8 Hz, H11), 5.24 (d, 2H, J = 12.0 Hz, Troc CH₂), 4.34-4.29 (m, 2H, OCH₂CH₂CH₂O), 4.20-4.11 (m, 4H, Troc CH₂ and OCH₂CH₂CH₂O), 4.00-3.96 (m, 8H, H11a and OCH₃ x

2), 3.84 (s, 6H, $\text{OCH}_3 \times 2$), 3.36 (dd, 2H, $J = 10.8, 16.6$ Hz, H1), 2.85 (d, 2H, $J = 16.5$ Hz, H1), 2.48–2.45 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 0.93 (s, 18H, TBS $\text{CH}_3 \times 6$), 0.30 and 0.27 (s $\times 2$, 12H, TBS $\text{CH}_3 \times 4$); ^{13}C NMR (100.6 MHz, CDCl_3) δ 162.5 (C_{quat}), 161.3 (C_{quat}), 159.2 (C_{quat}), 151.1 (C_{quat}), 148.1 (C_{quat}), 140.3 (C_{quat}), 126.2 (C13), 126.0 (C_{quat}), 123.2 (C_{quat}), 121.9 (C3), 119.3 (C_{quat}), 114.3 (C6), 111.9 (C14), 111.2 (C9), 95.2 (Troc CCl_3), 87.3 (C11), 74.8 (Troc CH_2), 65.0 ($\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 61.5 (C11a), 56.1 and 55.3 (OCH_3), 35.3 (C1), 28.8 ($\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 25.7 (TBS CH_3), 17.9 (TBS C_{quat}); MS (ES), m/z (relative intensity) 1357 (M^+ , 63), 1114 (48), 955 (59), 919 (78).

1,1'-[[(Propane-1,3-diyl)dioxy]bis[(11aS)-7-methoxy-2-(p-methoxybenzene)-1,11a-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]] (ZC-207)




10% Cd/Pd couple (461 mg, 3.73 mmol, 20 eq) was added to a rapidly stirring mixture of 15 (253 mg, 0.19 mmol), THF (5 mL) and 1 N NH_4OAc (5 mL). The reaction mixture was allowed to stir for 1.5 h when TLC showed the complete consumption of the starting material. The solids were filtered and rinsed with H_2O and DCM. The aqueous layer was extracted with DCM (3 \times 30 mL) and the organic extracts were combined, washed with brine (50 mL) and dried (MgSO_4). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (99.9:0.1 v/v $\text{CHCl}_3/\text{MeOH}$ then gradient to 95:5 v/v $\text{CHCl}_3/\text{MeOH}$) to afford ZC-207 as a yellow glass (132 mg, 96%): $[\alpha]^{20}_{\text{D}} = +880.0^\circ$ ($c = 0.22$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.79 (d, 2H, $J = 3.9$ Hz, H11), 7.44 (s, 2H, H6), 7.30 (s, 2H, H3), 7.24 (d, 4H, $J = 8.7$ Hz, H13), 6.81 (d, 4H, $J = 8.7$ Hz, H14), 6.79 (s, 2H, H9), 4.30–4.18 (m, 6H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$ and H11a), 3.86 (s, 6H, $\text{OCH}_3 \times 2$), 3.74 (s, 6H, $\text{OCH}_3 \times 2$), 3.48 (dd, 2H, $J = 11.8, 16.2$ Hz, H1), 2.85 (d, 2H, $J = 16.2$ Hz, H1), 2.38–2.32 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$); ^{13}C NMR (62.9 MHz, CDCl_3) δ

162.5 (C11), 161.3 (C_{quat}), 159.2 (C_{quat}), 151.1 (C_{quat}), 148.1 (C_{quat}),
 140.3 (C_{quat}), 126.2 (C13), 126.0 (C_{quat}), 123.2 (C_{quat}), 121.9 (C3),
 114.3 (C14), 111.9 (C9), 111.2 (C6), 65.4 (OCH₂CH₂CH₂O), 56.2 and
 55.3 (OCH₃), 53.8 (C11a), 35.6 (C1), 28.9 (OCH₂CH₂CH₂O); MS (ES),
 5 m/z (relative intensity) 741 (M⁺, 43), 660 (71).

Example 4 - 1,1'-[[(Propane-1,3-diyl)dioxy]bis[(11aS)-7-methoxy-2-(1-propenyl)-1,11a-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]] (ZC-211)


10 1,1'-[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butyldimethylsilyloxy)-7-methoxy-2-(1-propenyl)-1,10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]] (16)



15 A solution of TEA (0.11 mL, 81 mg, 0.80 mmol, 6.0 eq) in H₂O (3 mL) and EtOH (10 mL) was added to a solution of triflate 13 (192 mg, 0.13 mmol) in toluene (5 mL) at room temperature. To this mixture *trans*-propenylboronic acid (30 mg, 0.35 mmol, 2.6 eq) and Pd(PPh₃)₄ (6 mg, 5 μmol, 0.04 eq) were added. The reaction mixture was
 20 heated at 76 °C for 2 h when TLC (50:50 v/v EtOAc/hexane) revealed the complete consumption of the starting material. Excess solvent was removed and the residue was dissolved in EtOAc (15 mL), washed with H₂O (10 mL), brine (10 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was
 25 purified by flash column chromatography (80:20 v/v hexane/ EtOAc) to afford 16 as a light yellow glass (40 mg, 25%): [α]_D²⁰ = +75.0 ° (c = 0.20, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.28 (s, 2H, H6), 6.90 (s, 2H, H3), 6.78 (s, 2H, H9), 6.26 (d, 2H, J = 14.8 Hz, H12), 5.85 (d, 2H, J = 8.8 Hz, H11), 5.54 (dq, J = 6.8, 15.4 Hz, 2H, H13),
 30 5.23 (d, 2H, J = 12.0 Hz, Troc CH₂), 4.32-4.26 (m, 2H, OCH₂CH₂CH₂O), 4.18-4.11 (m, 4H, Troc CH₂ and OCH₂CH₂CH₂O), 3.94 (s, 6H, OCH₃ x 2), 3.89-3.83 (m, 2H, H11a), 3.07 (dd, 2H, J = 10.6, 15.9 Hz, H1), 2.60

(d, 2H, $J = 16.3$ Hz, H1), 2.46-2.43 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 1.85 (d, 6H, $J = 6.6$ Hz, H14), 0.90 (s, 18H, TBS CH_3 x 6), 0.28 and 0.25 (s x 2, 12H, TBS CH_3 x 4); ^{13}C NMR (100.6 MHz, CDCl_3) δ 163.6 (C_{quat}), 153.6 (C_{quat}), 150.4 (C_{quat}), 149.2 (C_{quat}), 127.7 (C_{quat}), 126.4 (C13), 125.6 (C_{quat}), 124.7 (C12 and C3), 123.5 (C_{quat}), 114.0 (C9), 110.7 (C6), 95.2 (Troc CCl_3), 87.2 (C11), 74.7 (Troc CH_2), 65.0 ($\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 61.4 (C11a), 56.1 (OCH_3), 33.9 (C1), 28.8 ($\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 25.6 (TBS CH_3), 18.4 (C14), 17.9 (TBS C_{quat}); MS (ES), m/z (relative intensity) 1246 ($[\text{M} + 22]^+$, 100), 1226 ($[\text{M} + 2]^+$, 88).

1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11aS)-7-methoxy-2-(1-propenyl)-1,11a-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]]] (ZC-211)




10% Cd/Pd couple (81 mg, 0.65 mmol, 20 eq) was added to a rapidly stirring mixture of 16 (40 mg, 0.03 mmol), THF (1 mL) and 1 N NH_4OAc (1 mL). The reaction mixture was allowed to stir for 1 h when TLC showed the complete consumption of the starting material. The solids were filtered and rinsed with H_2O and CHCl_3 . The aqueous layer was extracted with CHCl_3 (3 x 5 mL) and the organic extracts were combined, washed with H_2O (10 mL), brine (10 mL) and dried (MgSO_4). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (99:8:0.2 v/v $\text{CHCl}_3/\text{MeOH}$ then gradient to 97:3 v/v $\text{CHCl}_3/\text{MeOH}$) to afford ZC-211 as a yellow glass (14.7 mg, 74%): $[\alpha]^{20}_{\text{D}} = +1102.0^\circ$ ($c = 0.15$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.74 (d, 2H, $J = 3.8$ Hz, H11), 7.42 (s, 2H, H6), 6.84 (s, 2H, H3), 6.77 (s, 2H, H9), 6.19 (d, 2H, $J = 15.5$ Hz, H12), 5.52 (dq, $J = 6.8, 15.4$ Hz, 2H, H13), 4.27-4.16 (m, 6H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$ and H11a), 3.84 (s, 6H, OCH_3 x 2), 3.22 (dd, 2H, $J = 11.5, 16.1$ Hz, H1), 3.03 (dd, 2H, $J = 4.8, 16.2$ Hz, H1), 2.38-2.32 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 1.77 (d, 6H, $J = 6.6$ Hz, H14); ^{13}C NMR (100.6 MHz, CDCl_3) δ 162.6 (C11), 161.2 (C_{quat}), 151.1 (C_{quat}), 148.1

(C_{quat}), 140.3 (C_{quat}), 127.7 (C_{quat}), 126.9 (C13), 124.4 (C12 and C3),
 123.9 (C_{quat}), 119.3 (C_{quat}), 111.9 (C6), 111.2 (C9), 65.4
 (OCH₂CH₂CH₂O), 56.2 (OCH₃), 53.8 (C11a), 34.2 (C1), 28.8
 (OCH₂CH₂CH₂O), 18.5 (C14); MS (ES), *m/z* (relative intensity) 609
 (M⁺, 100).

Example 5 - 1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11aS)-7-methoxy-2-ethynylphenyl-1,11a-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]] (ZC-209)


1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11S,11aS)-10-(2,2,2-trichloroethoxycarbonyl)-11-(tert-butyldimethylsilyloxy)-7-methoxy-2-ethynylphenyl-1,10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-one]] (17)



A catalytic amount of Pd(PPh₃)₄ was added to a stirred mixture of triflate 13 (193 mg, 0.13 mmol), LiCl (34 mg, 0.80 mmol, 6.0 eq) and tributyl(phenylethynyl)tin (0.14 mL, 157 mg, 0.40 mmol, 3.0 eq) in dry THF (5 mL). The reaction mixture was heated at reflux for 2.5 h when TLC showed the complete consumption of the starting material. After cooling to room temperature, excess solvent was removed and the residue was dissolved in DCM (20 mL), followed by washin with 10 % NH₄OH (20 mL). The aqueous layer was extracted with DCM (3 x 20 mL), and the organic extracts were combined, washed with brine (50 mL) and dried (MgSO₄). Filtration and evaporation of solvent afforded the crude product which was purified by flash column chromatography (80:20 v/v hexane/EtOAc) to afford 17 as a yellow glass (162 mg, 90%): ¹H NMR (400 MHz, CDCl₃) δ 7.40-7.37 (m, 4H, H15), 7.26-7.19 (m, 10H, H3, H6, H16 and H17), 6.70 (s, 2H, H9), 5.85 (d, 2H, *J* = 8.8 Hz, H11), 5.15 (d, 2H, *J* = 12.0 Hz, Troc CH₂), 4.24-4.19 (m, 2H, OCH₂CH₂CH₂O), 4.12-4.02 (m,

4H, Troc CH₂ and OCH₂CH₂CH₂O), 3.86-3.79 (m, 8H, OCH₃ x 2 and H11a),
 3.15 (dd, 2H, J = 10.8, 16.5 Hz, H1), 2.63 (d, 2H, J = 16.5 Hz,
 H1), 2.37-2.35 (m, 2H, OCH₂CH₂CH₂O), 0.82 (s, 18H, TBS CH₃ x 6),
 0.22 and 0.18 (s x 2, 12H, TBS CH₃ x 4); ¹³C NMR (100.6 MHz, CDCl₃)
 5 δ 163.8 (C_{quat}), 153.6 (C_{quat}), 150.8 (C_{quat}), 149.3 (C_{quat}), 133.3 (C3),
 131.4 (C15), 128.4 (C16 and C17), 127.8 (C_{quat}), 125.1 (C_{quat}), 123.0
 (C_{quat}), 114.0 (C6), 110.0 (C9), 104.8 (C_{quat}), 95.2 (Troc CCl₃), 93.7
 (Calkyne), 86.9 (C11), 83.6 (Calkyne), 74.8 (Troc CH₂), 66.0
 (OCH₂CH₂CH₂O), 61.4 (C11a), 56.2 (OCH₃), 37.8 (C1), 28.8
 (OCH₂CH₂CH₂O), 25.6 (TBS CH₃), 17.8 (TBS C_{quat}); MS (ES), m/z
 0 (relative intensity) 1344 (M⁺, 8), 625 (100).

1,1'-[[[(Propane-1,3-diyl)dioxy]bis[(11aS)-7-methoxy-2-
 ethynylphenyl-1,11a-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepine-5-
 5 one]] (ZC-209)




10% Cd/Pd couple (314 mg, 2.55 mmol, 20 eq) was added to a rapidly
 stirring mixture of 17 (162 mg, 0.13 mmol), THF (4 mL) and 1 N
 NH₄OAc (4 mL). The reaction mixture was allowed to stir for 45 min
 20 when TLC showed the complete consumption of the starting material.
 The solids were filtered and rinsed with H₂O and CHCl₃. The aqueous
 layer was extracted with CHCl₃ (3 x 25 mL) and the organic extracts
 were combined, washed with H₂O (50 mL), brine (50 mL) and dried
 (MgSO₄). Filtration and evaporation of solvent afforded the crude
 25 product which was purified by flash column chromatography (99.8:0.2
 v/v CHCl₃/MeOH then gradient to 97.5:2.5 v/v CHCl₃/MeOH) to afford
 ZC-209 as a yellow glass (33 mg, 38%): ¹H NMR (400 MHz, CDCl₃) δ
 7.79 (d, 2H, J = 3.9 Hz, H11), 7.42 (s, 2H, H6), 7.39-7.37 (m, 4H,
 H15), 7.26-7.19 (m, 8H, H3, H16 and H17), 6.78 (s, 2H, H9), 4.30-
 30 4.19 (m, 6H, OCH₂CH₂CH₂O and H11a), 3.86 (s, 6H, OCH₃ x 2), 3.36

(dd, 2H, $J = 11.7, 16.4$ Hz, H1), 3.18 (dd, 2H, $J = 5.4, 16.4$ Hz, H1), 2.37-2.34 (m, 2H, $\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$); ^{13}C NMR (100.6 MHz, CDCl_3) δ 162.0 (C11), 161.3 (C_{quat}), 151.4 (C_{quat}), 148.2 (C_{quat}), 140.3 (C_{quat}), 132.8 (C3), 131.4 (C15), 128.4 (C16 and C17), 122.9 (C_{quat}), 118.7 (C_{quat}), 112.0 (C6), 111.2 (C9), 105.1 (C_{quat}), 94.3 (C_{alkyne}), 83.3 (C_{alkyne}), 65.5 ($\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$), 56.2 (OCH_3), 53.8 (C11a), 37.9 (C1), 28.8 ($\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}$); MS (ES), m/z (relative intensity) 729 (M^+ , 100).

Example 6 - Formation of key monomer intermediate ((1*S*,11*aS*)-7,8-Dimethoxy-11-(*tert*-butyldimethylsilyloxymethyl)-10-(2,2,2-trichloroethoxycarbonyl)-2-[[*(trifluoromethyl)sulfonyl*oxy]-1,2,3,10,11,11*a*-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one (27))


(2*S*)(4*R*)-*N*-(4,5-Dimethoxy-2-nitrobenzoyl)-2-(*tert*-butyldimethylsilyloxymethyl)-4-oxyacetylpyrrolidine (19)



A stirred solution of 6-nitroveratric acid (18) (1.49 g, 6.58 mmol; 1.2 Equiv.) in dry DMF (2 mL) and DCM (80 mL) was treated with *N*-(3-Dimethylaminopropyl)-*N*-ethyl carbodiimide hydrochloride (1.26 g, 6.58 mmol, 1.2 Equiv.) and cooled to 0°C. 1-Hydroxybenzotriazole hydrate (1.03 g, 6.58 mmol, 1.2 Equiv.) was added in portions and the reaction mixture was allowed to warm to room temperature. After stirring for 1 h, a solution of 3 (1.5 g, 5.48 mmol, 1.0 Equiv.) in DCM (80 mL) was added dropwise and the mixture stirred for a further 15 h. This was followed by heating at 55°C for a further 3 h. The reaction mixture was washed with NH_4Cl (60 mL), NaHCO_3 (60 mL), brine (60 mL), dried (MgSO_4) and concentrated *in vacuo*. Purification by flash chromatography (EtOAc/petroleum ether 4:6) afforded the product 19 as pale crystals (2.11g, 79.9%): m.p. 134°C; $[\alpha]_D^{19} = -91.9^\circ$ ($c = 0.408$, CHCl_3); ^1H NMR (CDCl_3 , 400 MHz) δ 0.12 (s, 6 H, $\text{Si}(\text{CH}_3)_2$), 0.93 (s, 9 H, $\text{C}(\text{CH}_3)_3$), 2.04 (s, 3

H, C=OCH₃), 2.20 - 2.28 (m, 1 H, H1a), 2.39 - 2.45 (m, 1 H, H1b), 3.22 (d, 1 H, *J* = 11.74 Hz, H3a), 3.47 (dd, 1 H, *J* = 11.89, 4.70 Hz, H3b), 3.75 (d, 1 H, *J* = 8.39 Hz, H11), 3.94 (s, 3 H, CH₃O7), 3.98 (s, 3 H, CH₃O8), 4.21 (d, 1 H, *J* = 9.76 Hz, H11), 4.56 (m, 1 H, H11a), 5.19 (m, 1 H, H2b), 6.73 (s, 1 H, H6), 7.68 (s, 1 H, H9); ¹³C NMR (CDCl₃, 100 MHz) δ 171.0 (C=O), 166.3 (C5), 154.0 (C7), 149.1 (C8), 137.6 (C5-6), 127.8 (C9-10), 109.1 (C6), 107.3 (C9), 72.9 (C2), 62.6 (C11), 57.4 (C11a), 56.5, 56.6 (CH₃O7 and CH₃O8), 52.0 (C3), 33.0 (C1), 25.7 (C(CH₃)₃), 21.2 (C=OCH₃), 18.1 (Cquat), -5.5 (SiCH₃)₂; IR (film) 2953, 2856, 1740 (C=O), 1648 (C=O), 1579, 1525, 1463, 1427, 1338, 1278, 1243, 1225, 1115, 1070, 1028, 1004, 837, 780 cm⁻¹; MS (EI) *m/z* (relative intensity) 483 ([M + H]⁺, 100); Elem. Anal. calculated for C₂₂H₃₄N₂O₈Si: C, 54.75; H, 7.10; N, 5.80. Found: C, 54.50; H, 7.08; N, 5.79.


(2*S*,4*R*)-*N*-(2-Amino-4,5-dimethoxybenzoyl)-2-(*tert*-butyldimethylsilyloxymethyl)-4-oxyacetylpyrrolidine (20)



A suspension of 19 (61.75 g, 127.9 mmol; 1 Equiv.) and 10% w/w Pd/C (6.17 g) in ethanol (400 mL) was agitated under a H₂ atmosphere (45 psi) using Parr apparatus over a period of 3 h. The suspension was filtered through celite and the filtrate was evaporated *in vacuo* to afford 20 as a pale yellow oil (57.5 g, 99%): [α]_D¹⁹ = -105.1° (c = 0.490, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ 0.03 (s, 6 H, Si(CH₃)₂), 0.88 (s, 9 H, C(CH₃)₃), 1.98 (s, 3 H, C=OCH₃), 2.09 - 2.15 (m, 1 H, H1a), 2.32-2.39 (m, 1 H, H1b), 3.57-3.64 (m, 2 H, H3a and H11), 3.73-3.77 (m, 4 H, H3b and CH₃O8), 3.82 (s, 3 H, CH₃O7), 4.00-4.11 (m (br), 1 H, H11), 4.40-4.60 (m, 1 H, H11a), 5.21-5.27 (m, 1 H, H2b), 6.21 (s, 1 H, H9), 6.69 (s, 1 H, H6); ¹³C NMR (CDCl₃, 100 MHz) δ 170.6 (C=O), 170.0 (C5), 151.9 (C7), 141.7 (C9-10), 140.9 (C8), 112.2 (C6), 110.9 (C5-6), 100.6 (C9), 73.5 (C2), 62.6 (C11), 57.0 (C11a), 55.8, (CH₃O7), 56.6 (CH₃O8), 56.3 (C3), 32.9 (C1), 25.8 (C(CH₃)₃), 21.1 (C=OCH₃), 18.1 (Cquat), -5.5 (SiCH₃)₂; IR (film)


3455, 3355 (NH₂), 2931, 2857, 1740 (C=O), 1628 (C=O), 1594, 1516, 1401, 1238, 1165, 1120, 1006, 837, 778, 668 cm⁻¹; MS (EI) *m/z* (relative intensity) 453 ([M + H]⁺, 100).

- 5 (2*S*, 4*R*)-*N*-(4,5-Dimethoxy-2-(2,2,2-trichloroethoxycarbonylamino)-benzoyl)-2-(*tert*-butyldimethylsilyloxymethyl)-4-oxyacetylpyrrolidine (**21**)



- 2,2,2-Trichloroethyl chloroformate (1.61 mL, 11.68 mmol, 2.2
 10 Equiv.) was added to a stirred solution of **20** (2.40 g, 5.31 mmol; 1.0 Equiv.) and pyridine (1.72 mL, 21.24 mmol, 4.0 Equiv.) in dry DCM (70 mL) at -20°C. The mixture was allowed to warm to room temperature and stirred for a further 2.5 h. The reaction mixture was washed with NH₄Cl (2 x 60 mL), CuSO₄ (60 mL), water (60 mL),
 15 brine (60 mL), dried (MgSO₄), filtered and concentrated *in vacuo* to afford **21** as a colourless oil (3.32 g, 99 %): [α]_D²⁰ = -43.4° (c = 0.980, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ 0.28 (s, 6 H, Si(CH₃)₂), 0.87 (s, 9 H, C(CH₃)₃), 1.96 (s, 3 H, C=OCH₃), 2.10 - 2.16 (m, 1 H, H1a), 2.35 - 2.41 (m, 1 H, H1b), 3.55 - 3.65 (br m, 1 H, H3a), 3.68 - 3.74 (m, 2 H, H11 and H3b), 3.78 (s, 3 H, CH₃O7), 3.91 (s, 3 H, CH₃O8), 4.06 - 4.10 (m, 2 H, Cl₃CCH₂ rotamers and H11), 4.55 - 4.65 (m, 1 H, H11a), 4.74 - 4.82 (m, 1 H, Cl₃CCH₂, rotamers), 5.20 - 5.25 (m, 1 H, H2b), 6.75 (s, 1 H, H6), 7.82 (br s, 1 H, H9), 9.38 (br s, 1 H, NH); ¹³C NMR (CDCl₃, 100 MHz) δ 170.5 (C=O), 169.2 (C5), 153.1,
 20 - 3.74 (m, 2 H, H11 and H3b), 3.78 (s, 3 H, CH₃O7), 3.91 (s, 3 H, CH₃O8), 4.06 - 4.10 (m, 2 H, Cl₃CCH₂ rotamers and H11), 4.55 - 4.65 (m, 1 H, H11a), 4.74 - 4.82 (m, 1 H, Cl₃CCH₂, rotamers), 5.20 - 5.25 (m, 1 H, H2b), 6.75 (s, 1 H, H6), 7.82 (br s, 1 H, H9), 9.38 (br s, 1 H, NH); ¹³C NMR (CDCl₃, 100 MHz) δ 170.5 (C=O), 169.2 (C5), 153.1,
 25 151.4 (OC=ONH, rotamers), 151.9 (C8), 144.1 (C7), 132.2 (C5-6), 114.8 (C9-10), 111.1 (C6), 104.3 (C9), 95.3, 93.9 (Cl₃C, rotamers), 77.3, 76.3 (Cl₃CCH₂, rotamers), 73.4 (C2), 62.3 (C11), 57.3 (C11a), 57.2 (C3), 56.1, 56.2 (CH₃O7 and CH₃O8), 32.5 (C1), 25.8 (C(CH₃)₃), 21.1 (C=OCH₃), 18.1 (Cquat), -5.5 (SiCH₃)₂; IR (film) 3318, 2954,
 30 2858, 1774, 1743 (C=O and OC=ON), 1601 (NC=O), 1525, 1464, 1422, 1398, 1230, 1200, 1125, 1004, 836, 777, 720 cm⁻¹; MS (EI) *m/z* (relative intensity) 629 ([M + H]⁺, 100).


(2*S*, 4*R*)-*N*-(4,5-Dimethoxy-2-(2,2,2-trichloroethoxycarbonylamino)-benzoyl)-2-hydroxymethyl-4-oxyacetylpyrrolidine (**22**)



A mixture of acetic acid (81 mL) and water (18 mL) was added to a stirred solution of **21** (5.10 g, 8.12 mmol; 1.0 Equiv.) in THF (45 mL). The resulting solution was stirred at room temperature for a period of 48 h. The THF was removed *in vacuo* and the resulting mixture was neutralised to pH 7 with solid NaHCO₃ (Caution! vigorous effervescence). The resulting aqueous layer was extracted with DCM (5 x 150 mL), dried (MgSO₄) and concentrated *in vacuo*. The oily residue was subjected to flash chromatography (EtOAc/Petroleum ether 6:4) to give **22** as a foam (4.16 g, 99%):


$[\alpha]_D^{18} = -65.0^\circ$ ($c = 0.500$, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ 2.01 (s, 3 H, C=OCH₃), 2.05 - 2.13 (m, 1 H, H1b), 2.27 (dd, 1 H, $J = 14.20, 7.62$ Hz, H1a), 3.62 (d, 1 H, $J = 12.36$ Hz, H3a), 3.65 - 3.74 (m, 1 H, H11), 3.76 (dd, 1 H, $J = 12.57, 3.80$ Hz, H3b), 3.85 (s, 3 H, CH₃O7), 3.93 (s, 3 H, CH₃O8), 3.98 - 4.08 (m, 1 H, H11), 4.55 - 4.65 (m, 1 H, H11a), 4.79 (d, 1 H, $J = 12.09$ Hz, Cl₃CCH₂), 4.84 (d, 1 H, $J = 12.04$ Hz, Cl₃CCH₂), 5.18 - 5.25 (m, 1 H, H2b), 6.80 (s, 1 H, H6), 7.73 (br s, 1 H, H9), 9.03 (br s, 1 H, NH); ¹³C NMR (CDCl₃, 100 MHz) δ 170.6 (C=O), 170.4 (C5), 152.1 (OC=ONH), 151.5 (C8), 144.6 (C7), 131.2 (C5-6), 115.5 (C9-10), 110.8 (C6), 104.8 (C9), 95.3 (Cl₃C), 74.4 (Cl₃CCH₂), 72.5 (C2), 64.5 (C11), 58.8 (C11a), 56.6 (C3), 56.5 (CH₃O7), 56.1 (CH₃O8), 33.7 (C1), 21.1 (C=OCH₃); IR (film) 3358 (br NH and OH), 3015, 2941, 1740 (C=O and OC=ON), 1602 (NC=O), 1525, 1463, 1433, 1398, 1231, 1174, 1125, 1037, 969, 817, 756 cm⁻¹; MS (EI) m/z (relative intensity) 363 ([M - Cl₃CCH₂O]⁺, 100), 513 ([M + H]⁺, 95).

(11*S*,11*aS*,2*R*)-7,8-Dimethoxy-11-hydroxy-2-oxyacetyl-10-(2,2,2-trichloroethoxycarbonyl)-1,2,3,10,11,11*a*-hexahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one (23)



- 5 Diodobenzene diacetate (83.7 g, 259.8 mmol, 1.78 Equiv.) and 2,2,6,6-tetramethylpiperidine nitroxyl (TEMPO) (4.50 g, 28.8 mmol, 0.2 Equiv.) were added to a stirred solution of 22 (74.7 g, 145.4 mmol; 1.0 Equiv.) in dry DCM (1.5 L). The reaction mixture was stirred at room temperature for 15 h and diluted with DCM (500 mL).
- 10 The organic phase was washed with a sat^d sodium bisulphite (700 mL) and the aqueous layer was back-extracted with DCM (3 x 200 mL). The organic layers were combined, dried (MgSO₄), filtered and concentrated *in vacuo*. Purification by flash chromatography (EtOAc/Petroleum ether 7:3) afforded 23 as a white glass (57.65 g, 77 %): $[\alpha]_D^{18} = +99.4^\circ$ ($c = 0.483$, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ 2.04 (s, 3 H, C=OCH₃), 2.34 - 2.46 (m, 2 H, H1b and H1a), 3.69 - 3.77 (m, 2 H, H3b and 11a), 3.92 (s, 3 H, CH₃O8), 3.95 (s, 3 H, CH₃O7), 4.05 (dd, 1 H, $J = 13.13, 2.35$ Hz, H3a), 4.23 (d, 1 H, $J = 12.02$ Hz, Cl₃CCH₂), 5.25 (d, 1 H, $J = 12.02$ Hz, Cl₃CCH₂), 5.38 (p, 1 H, $J = 4.10$ Hz, H2b), 5.70 (dd, 1 H, $J = 9.66, 3.79$ Hz, H11), 6.81 (s, 1 H, H9), 7.27 (s, 1 H, H6); ¹³C NMR (CDCl₃, 100 MHz) δ 170.3 (C=O), 167.4 (C5), 154.4 (OC=ONH), 151.1 (C8), 148.8 (C7), 127.5 (C9-10), 124.7 (C5-6), 112.7 (C9), 110.7 (C6), 95.0 (Cl₃C), 87.6 (C11), 75.0 (Cl₃CCH₂), 71.4 (C2), 58.3 (C11a), 56.19, 56.13 (CH₃O8 and CH₃O7), 51.1 (C3), 35.9 (C1), 21.0 (C=OCH₃); IR (film) 3421 (br OH), 3008, 2946, 1719 (C=O), 1623 (OC=ON), 1603 (NC=O), 1515, 1429, 1374, 1304, 1238, 1212, 1057, 870, 819, 758, 711, 644 cm⁻¹; MS (EI) m/z (relative intensity) 511 ($[M - H]^+$, 100), 512.5 ($[M + H]^+$, 99).


(11*S*,11*aS*,2*R*)-7,8-Dimethoxy-2-oxyacetyl-11-(tert-butyltrimethylsilyloxymethyl)-10-(2,2,2-trichloroethoxycarbonyl)-1,2,3,10,11,11*a*-hexahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one (24)



5
10
15
20
25
30


TBDMSOTf (0.56 mL, 2.44 mmol; 1.5 Equiv.) was added to a stirred solution of 23 (834 mg, 1.63 mmol, 1.0 Equiv.) and 2,6-lutidine (0.38 mL, 3.26 mmol, 2.0 Equiv.) in dry DCM (5 mL). The mixture was stirred at room temperature for 30 min and diluted with DCM (20 mL). The mixture was washed with sat^d CuSO₄ (2 x 20 mL), sat^d NaHCO₃ (30 mL), brine (30 mL), dried (MgSO₄) and concentrated *in vacuo* to afford 6 as white glass (1.01 g, 99 %): $[\alpha]_D^{18} = +52.7^\circ$ ($c = 0.237$, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ 0.23 (s, 6 H, Si(CH₃)₂), 0.87 (s, 9 H, C(CH₃)₃), 2.03 (s, 3 H, C=OCH₃), 2.18 - 2.25 (m, 1 H, H1b), 2.30 - 2.40 (m, 1 H, H1a), 3.66 - 3.72 (m, 2 H, H3b and 11a), 3.89 (s, 3 H, CH₃O8), 3.95 (s, 3 H, CH₃O7), 4.13 (d, 1 H, $J = 13.40$ Hz, H3a), 4.18 (d, 1 H, $J = 12.04$ Hz, Cl₃CCH₂), 5.23 (d, 1 H, $J = 12.03$ Hz, Cl₃CCH₂), 5.37 (p, 1 H, $J = 2.62$ Hz, H2b), 5.77 (d, 1 H, $J = 8.95$ Hz, H11), 6.74 (s, 1 H, H9), 7.28 (s, 1 H, H6); ¹³C NMR (CDCl₃, 100 MHz) δ 170.3 (C=O), 167.9 (C5), 153.5 (OC=ONH), 151.0 (C8), 148.9 (C7), 128.0 (C9-10), 125.4 (C5-6), 112.9 (C9), 110.6 (C6), 95.2 (Cl₃C), 88.2 (C11), 74.7 (Cl₃CCH₂), 71.7 (C2), 60.5 (C11a), 56.1 (CH₃O7), 56.9 (CH₃O8), 51.2 (C3), 36.2 (C1), 25.5 (C(CH₃)₃) 21.0 (C=OCH₃), 17.8 (Cquat), -4.3 and -5.3 (Si(CH₃)₂); IR (film) 3023, 2956, 1738 (C=O), 1718 (OC=ON), 1644 (NC=O), 1605, 1518, 1466, 1428, 1411, 1376, 1301, 1245, 1214, 1116, 1075, 1041, 1023, 842, 784, 756, 730, 712 cm⁻¹; MS (EI) m/z (relative intensity) 627 ([M + H]⁺, 100).

(11S,11aS,2R)-7,8-Dimethoxy-2-hydroxy-11-(tert-butyltrimethylsilyloxymethyl)-10-(2,2,2-trichloroethoxycarbonyl)-1,2,3,10,11,11a-hexahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one (25)




A solution of K_2CO_3 (732 mg, 5.30 mmol; 1.0 Equiv.) in water (15 mL) was added to a stirred solution of **24** (3.32 g, 5.30 mmol, 1.0 Equiv.) in MeOH (15 mL) and THF (5 mL). The mixture was stirred at room temperature for 5 h and then excess solvent was removed by rotary evaporation at reduced pressure. The remaining aqueous residue was neutralised to pH 7 with 1N HCl and extracted with EtOAc (4 x 30 mL). The organic layers were combined, washed with brine (50 mL), dried ($MgSO_4$), filtered and concentrated *in vacuo*. The resulting oil was subjected to flash chromatography (EtOAc) to afford **25** as a white glass (2.84 g, 92%): $[\alpha]_D^{22} = +58.3^\circ$ ($c = 0.587$, $CHCl_3$); 1H NMR ($CDCl_3$, 500 MHz) δ 0.22 (s, 6 H, $Si(CH_3)_2$), 0.87 (s, 9 H, $C(CH_3)_3$), 2.06 – 2.11 (m, 1 H, **H1b**), 2.28 – 2.33 (m, 1 H, **H1a**), 3.60 (dd, 1 H, $J = 4.31, 12.67$ Hz, **H3b**), 3.71 (q, 1 H, $J = 7.42, 15.67$ Hz, **H11a**), 3.88 (s, 6 H, CH_3O8 and CH_3O7), 4.01 (d, 1 H, $J = 12.93$ Hz, **H3a**), 4.17 (d, 1 H, $J = 12.03$ Hz, Cl_3CCH_2), 4.58 (br s, 1 H, **H2b**), 5.23 (d, 1 H, $J = 12.03$ Hz, Cl_3CCH_2), 5.54 (d, 1 H, $J = 9.00$ Hz, **H11**), 6.73 (s, 1 H, **H9**), 7.21 (s, 1 H, **H6**); ^{13}C NMR ($CDCl_3$, 125 MHz) δ 168.4 (**C5**), 153.6 ($OC=ONH$), 151.9 (**C8**), 148.8 (**C7**), 128.0 (**C9-10**), 125.5 (**C5-6**), 112.8 (**C9**), 110.5 (**C6**), 95.2 (Cl_3C), 88.2 (**C11**), 74.7 (Cl_3CCH_2), 69.5 (**C2**), 60.8 (**C11a**), 56.0, 55.9 (CH_3O8 and CH_3O7), 54.0 (**C3**), 38.8 (**C1**), 25.6 ($C(CH_3)_3$), 17.8 (**Cquat**), -4.2 and -5.2 ($Si(CH_3)_2$); IR (film) 3400 (br OH), 2932, 1731 ($OC=ON$), 1629 ($NC=O$), 1604, 1515, 1456, 1430, 1407, 1302, 1273, 1214, 1117, 1075, 987, 837, 754, 712, 638 cm^{-1} ; MS (EI) m/z (relative intensity) 585 ($[M + H]^+$, 100).

(11*S*,11*aS*)-7,8-Dimethoxy-2-oxo-11-(tert-butyltrimethylsilyloxymethyl)-10-(2,2,2-trichloroethoxycarbonyl)-1,2,3,10,11,11*a*-hexahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one (26)



Diodobenzene diacetate (2.44 g, 7.58 mmol, 1.78 Equiv.) and 2,2,6,6-tetramethylpiperidine nitroxyl (TEMPO) (133 mg, 0.85 mmol, 0.2 Equiv.) were added to a stirred solution of 25 (2.49 g, 4.26 mmol; 1.0 Equiv.) in dry DCM (40 mL). The mixture was stirred at room temperature for 18 h and the reaction mixture was diluted with DCM (50 mL). The organic phase was washed with sat^d sodium bisulphite (2 x 25 mL), brine (40 mL), dried (MgSO₄) and concentrated *in vacuo*. Purification by flash chromatography (EtOAc/Petroleum ether 6:4) afforded 26 as a white glass. (2.26 g, 91 %): $[\alpha]_D^{22} = +95.0^\circ$ ($c = 0.795$, CHCl₃); ¹H NMR (CDCl₃, 500 MHz) δ 0.22, 0.23 (two s, 6 H, Si(CH₃)₂), 0.86 (s, 9 H, C(CH₃)₃), 2.56 (dd, 1 H, $J = 3.12, 19.60$, H1b), 2.96 (dd, 1 H, $J = 10.29, 18.9$, H1a), 3.90 (s, 3 H, CH₃O8), 3.95 – 3.99 (m, 5 H, H3b, H11a and CH₃O7), 4.21 (d, 1 H, $J = 12.02$ Hz, Cl₃CCH₂), 4.32 (d, 1 H, $J = 20.93$ Hz, H3a), 5.24 (d, 1 H, $J = 12.03$ Hz, Cl₃CCH₂), 5.83 (d, 1 H, $J = 9.26$ Hz, H11), 6.77 (s, 1 H, H9), 7.25 (s, 1 H, H6); ¹³C NMR (CDCl₃, 125 MHz) δ 207.8 (C2), 168.0 (C5), 153.7 (OC=ONH), 151.4 (C8), 149.2 (C7), 128.0 (C9–10), 124.5 (C5–6), 113.0 (C9), 110.4 (C6), 95.1 (Cl₃C), 87.4 (C11), 74.8 (Cl₃CCH₂), 58.9 (C11a), 56.2 (CH₃O7), 56.0 (CH₃O8), 52.8 (C3), 40.3 (C1), 25.5 (C(CH₃)₃), 17.8 (Cquat), –4.2 and –5.3 (Si(CH₃)₂); IR (film) 2934, 1763 (C=O), 1720 (OC=ON), 1649 (NC=O), 1604, 1515, 1402, 1274, 1217, 1120, 1075, 1002, 866, 834, 756, 712 cm^{–1}; MS (EI) m/z (relative intensity) 615 ([M + MeOH]⁺, 100).

(11*S*,11*aS*)-7,8-Dimethoxy-11-(tert-butyldimethylsilyloxymethyl)-10-(2,2,2-trichloroethoxycarbonyl)-2-[[trifluoromethyl)sulfonyl]oxy]-1,2,3,10,11,11*a*-tetrahydro-5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-5-one (27)




Pyridine (1.71 mL; 21.17 mmol; 7 Equiv.) and trifluoromethanesulfonic anhydride (3.56 mL; 21.17 mmol; 7 Equiv) were added to a stirred solution of 27 (1.76 g, 3.02 mmol; 1 Equiv.) in dry CHCl₃ (50 mL) at 0-5 °C (ice bath). The ice bath was removed and the reaction mixture was stirred at room temperature for 3 h. At this point TLC analysis revealed the persistence of starting material. As a result, one extra equivalent of pyridine and Tf₂O was added (0.24 and 0.51 mL, respectively). The reaction was stirred for an additional 1 h until complete consumption of the starting material was observed by TLC. The reaction mixture was washed with water, sat^d CuSO₄, sat^d NaHCO₃, dried (MgSO₄), filtered and the solvent was evaporated *in vacuo*. The resulting residue was purified by flash chromatography (EtOAc/Petroleum ether 2:8) to give 27 as a pale yellow glass (1.12 g, 52%): [α]_D²² = +56.2° (c = 0.587, CHCl₃); ¹H NMR (CDCl₃, 500 MHz) δ 0.25, 0.27 (two s, 6 H, Si(CH₃)₂), 0.88 (s, 9 H, C(CH₃)₃), 2.82 (dd, 1 H, J = 2.79, 16.66 Hz, H1b), 3.33 (ddd, 1 H, J = 1.96, 10.73, 16.59 Hz, H1a), 3.90 (s, 3 H, CH₃O8), 3.94 (s, 3 H, CH₃O7), 3.88 - 3.96 (m, 1 H, H11a), 4.20 (d, 1 H, J = 12.00 Hz, Cl₃CCH₂), 5.23 (d, 1 H, J = 12.00 Hz, Cl₃CCH₂), 5.93 (d, 1 H, J = 9.26 Hz, H11), 6.74 (s, 1 H, H9), 7.17 (s, 1 H, H3), 7.23 (s, 1 H, H6); ¹³C NMR (CDCl₃, 125 MHz) δ 164.9 (C5), 153.6 (OC=ONH), 151.8 (C8), 149.3 (C7), 136.0 (C2), 127.9 (C9-10), 123.8 (C5-6), 121.0 (C3), 119.8 (CF₃), 113.2 (C9), 110.7 (C6), 95.1 (Cl₃C), 86.4 (C11), 74.9 (Cl₃CCH₂), 60.6 (C11a), 56.2 (CH₃O7), 56.0 (CH₃O8), 34.4 (C1), 25.5 (C(CH₃)₃), 17.8 (Cquat), -4.2 and -5.4 (Si(CH₃)₂); IR (film) 3008, 2930, 2858, 1725 (OC=ON), 1651 (NC=O), 1516, 1423, 1214, 1136,

1078, 897, 835, 783, 760, 713, 642 cm^{-1} ; MS (EI) m/z (relative intensity) 715 ($[M + H]^+$, 100).

Example 7 - (11aS)-7,8-Dimethoxy-2-(1-propenyl)-1,11a-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one (29)

(11S,11aS)-7,8-Dimethoxy-11-(tert-butyldimethylsilyloxymethyl)-10-(2,2,2-trichloroethoxycarbonyl)-2-(1-propenyl)-1,2,3,10,11,11a-tetrahydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one (28)




10 A mixture of Et_3N (0.34 mL, 2.48 mmol, 3 Equiv.), water (0.5 mL) and EtOH (1.70 mL) was added to a solution of **27** (590 mg, 0.826 mmol, 1 Equiv.) in toluene (1.70 mL) with vigorous stirring. The reaction mixture was treated with *trans*-propenylboronic acid (92.2 mg, 1.07 mmol, 1.3 Equiv.) and $\text{Pd}(\text{PPh}_3)_4$ (19 mg, 0.016 mmol, 0.02 Equiv.). After 2 h stirring at room temp TLC revealed no reaction.

15 The mixture was then heated to reflux (111°C) for 30 min after which time TLC showed the complete consumption of starting material. The reaction mixture was allowed to cool to room temperature and diluted with EtOAc (30 mL). The organic phase was

20 extracted with water (20 mL), brine (20 mL), dried (MgSO_4), filtered and the solvent was evaporated *in vacuo*. The resulting residue was purified by flash chromatography (EtOAc /Petroleum ether 2:8) to afford the product **10** (297 mg, 59%): $[\alpha]_D^{22} = +78.2^\circ$ ($c = 0.582$, CHCl_3); ^1H NMR (CDCl_3 , 500 MHz) δ 0.24, 0.26 (two s, 6 H, $\text{Si}(\text{CH}_3)_2$), 0.89 (s, 9 H, $\text{C}(\text{CH}_3)_3$), 1.83 (dd, 3 H, $J = 0.79$, 6.69 Hz, **H14**), 2.58 (dd, 1 H, $J = 3.11$, 16.45 Hz, **H1b**), 3.06 (dd, 1 H, $J = 10.59$, 16.40 Hz, **H1a**), 3.82 – 3.88 (m, 1 H, **H11a**), 3.89 (s, 3 H, **CH₃O8**), 3.94 (s, 3 H, **CH₃O7**), 4.17 (d, 1 H, $J = 12.04$ Hz, **Cl₃CCH₂**), 5.23 (d, 1 H, $J = 12.04$ Hz, **Cl₃CCH₂**), 5.50 – 5.56 (m, 1 H, **H13**), 5.84 (d, 1 H, $J = 8.86$ Hz, **H11**), 6.25 (d, 1 H, $J = 14.80$ Hz, **H12**), 6.75 (s, 1 H, **H9**), 6.88 (s, 1 H, **H3**), 7.24 (s, 1 H, **H6**); ^{13}C NMR (CDCl_3 , 125 MHz) δ 163.6 (**C5**), 153.6 (**OC=ONH**), 151.1 (**C8**), 149.0

(C7), 127.9 (C9-10), 126.5 (C13), 125.5 (C5-6), 124.7 (C12), 124.6 (C3), 123.6 (C2), 113.0 (C9), 110.5 (C6), 95.2 (Cl₃C), 87.2 (C11), 74.7 (Cl₃CCH₂), 61.4 (C11a), 56.2 (CH₃O7), 56.0 (CH₃O8), 33.9 (C1), 25.6 (C(CH₃)₃), 18.4 (C14), 17.9 (Cquat), -4.2 and -5.1 (Si(CH₃)₂);
5 IR (film) 2961, 2935, 2858, 1723 (OC=ON), 1648 (NC=O), 1514, 1404, 1272, 1216, 1076, 837, 756, 644 cm⁻¹; MS (EI) *m/z* (relative intensity) 607 ([M + H]⁺, 100).


10 (11aS)-7,8-Dimethoxy-2-(1-propenyl)-1,11a-dihydro-5H-pyrrolo[2,1-c][1,4]benzodiazepin-5-one (29)



10% Cd/Pd couple (418 mg, 3.38 mmol, 8.2 Equiv.) was added to a rapidly stirring mixture of 28 (250 mg, 0.412 mmol, 1 Equiv.), THF (3 mL) and 1 N ammonium acetate (3 mL). The reaction mixture was
15 allowed to stir for 3.5 h when TLC showed the complete consumption of the starting material. The solids were filtered and rinsed with H₂O and DCM. The aqueous layer was extracted with DCM (3 X 30 mL) and the organic extracts were combined, washed with brine (50 mL), dried (MgSO₄), filtered and evaporated in vacuo to provide the
20 crude product. Purification by flash column chromatography (0.5% → 2% MeOH in CHCl₃) afforded 29 as a yellow glass (96 mg, 78%):
[α]_D²⁴ = +989° (c = 0.890, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ 1.85 (d, 3 H, J = 6.26 Hz, H14), 3.12 (dd, 1 H, J = 4.95, 16.27 Hz, H1b), 3.30 (dd, 1 H, J = 11.80, 15.27 Hz, H1a), 3.93 (s, 3 H, CH₃O8),
25 3.95 (s, 3 H, CH₃O7), 4.30 (dt, J = 4.95, 15.70 Hz, H11a), 5.55 - 5.65 (m, 1 H, H13), 6.27 (d, J = 16.04 Hz, H12), 6.81 (s, 1 H, H9), 6.92 (s, 1 H, H3), 7.50 (s, 1 H, H6), 7.82 (d, 1 H, J = 4.00 Hz, H11); ¹³C NMR (CDCl₃, 100 MHz) δ 162.7 (C11), 161.2 (C5), 151.8 (C8), 147.7 (C7), 140.4 (C9-10), 126.9 (C13), 124.46, 124.43 (C12 and C3), 123.9 (C2), 119.2 (C5-6), 111.6 (C6), 109.8 (C9), 56.18, 56.11 (CH₃O7 and CH₃O8), 53.8 (C11a), 34.2 (C1), 18.4 (C14); IR (film) 3018, 2997, 2930, 2847, 1610, 1598 (C=O), 1507, 1431, 1344,

1254, 1214, 1101, 1065, 959, 862, 779 cm^{-1} ; MS (EI) m/z (relative intensity) 331 ($[\text{M} + \text{MeOH}]^+$, 100), 299 ($[\text{M} + \text{H}]^+$, 45).

Example 8 - Parallel Synthesis of C2-substituted PBDs (30a-al)



30a-al

(i) A solution of triethylamine (0.12 mL, 6 equiv.) in water (0.2 mL) and ethanol (1.0 mL) was poured into a stirred solution of 27 (100 mg, 0.14 mmol, 1 Equiv.) in toluene (1.0 mL) in an Emrys™ Process vial. To the solution was added the appropriate boronic acid (1.07 equiv.) and polystyrene triphenylphosphine palladium (0) PS-PPh₃Pd (15 mg, 0.0014 mmol, 0.01 Equiv). The vial was sealed with a Reseal™ septum, and the suspension then irradiated at 100°C for 20 minutes under microwave radiation using an Emrys™ Optimizer Microwave Station when TLC showed absence of starting material. N,N-diethanolaminomethyl polystyrene PS-DEAM (87.5 mg, 0.14 mmol, 1.0 equiv. [13 equiv. on boronic acid excess]) was added to the reaction, and the suspension was irradiated at 100°C for 10 minutes under microwave radiation (as above). Water (3 mL) was added to the reaction, and the suspension shaken for 10 minutes. The mixture was then transferred to a phase separator (PS) cartridge fitted with a selectively permeable frit pre-conditioned with CH₂Cl₂ (3 mL) and coupled to a Na₂SO₄ cartridge. Extraction with CH₂Cl₂ (3 x 5 mL), followed by concentration *in vacuo* yielded an oil which was redissolved in THF (1.5 mL) and 1N ammonium acetate (1.5 mL).

(ii) To the mixture resulting from the previous step was added 10% Cd/Pb (140 mg, 1.12 mmol, 8.0 Equiv.) while vigorously stirring and the reaction was kept for 1 hour at room temperature when TLC showed absence of starting material. The mixture was poured into an identical PS cartridge pre-conditioned with CH₂Cl₂ (3 mL) and coupled to a Na₂SO₄ cartridge. Extraction with CH₂Cl₂ (3 x 5 mL) followed by concentration *in vacuo* gave an oil which was subjected

to flash chromatography (EtOAc:Hexane 1:1) afforded the desired compounds, which are listed below.

Compound	R''	Yield (%) [†]	[α] _D ²⁵
30a	Phenyl	62	+1012°
30b	4-methylphenyl	39	+916°
30c	2-methylphenyl	38	+802°
30d	4-ethylphenyl	45	+985°
30e	2,6-dimethylphenyl	35	+549°
30f	4-methoxyphenyl	99 [‡]	+976°
30g	3-methoxyphenyl	---	+825°
30h	4-tert-butylphenyl	35	+781°
30i	4-fluorophenyl	41	+978°
30j	4-chlorophenyl	67	+796°
30k	4-biphenyl	44	+792°
30l	4-phenoxyphenyl	47	+823°
30m	2-naphthyl	47	+720°
30n	3,4-methylenedioxyphenyl	46	+837°
30o	trans-2-(4-methylphenyl)vinyl	58	+990°
30p	2-thiophenyl	59	+974°
30q	trans-propenyl	78 [‡]	+989°
30r	4-dimethylaminophenyl	44	+858°
30s	4-methylthiophenyl	18	+979°
30t	4-vinylphenyl	36	+796°
30u	3,4-dichlorophenyl	41	+641°
30v	4-trifluoromethylphenyl	28	+785°
30w	4-isopropylphenyl	44	+985°
30x	4-cyanophenyl	42	+1000°
30y	3-pyridinyl	14	+1200°
30z	4-pyridinyl	27	+859°
30aa	4-formylphenyl	42	+928°
30ab	4-carboxylphenyl	43	---
30ac	2,6-dimethoxyphenyl	18.5	---
30ad	4-acetanilide	26.3	---
30ae	4-aminophenyl	21.3	---
30af	1-naphthyl	32	---
30ag	5-indole	34	---
30ah	3-aminophenyl	41	---
30ai	2,6-difluorophenyl	14	---
30aj	1-pyrenyl	39	---
30ak	4-hydroxyphenyl	14	---
30al	trans-hexenyl	---	---

5

[†] Overall yield, including Suzuki coupling followed by imine formation.

[‡] yield calculated from Troc-deprotection final step only.

*Concentration range: 0.047-0.89 g/100mL; Temp.: 20-29°C. All samples were dissolved in HPLC grade chloroform stabilised with amylene, purchased from Fisher Chemicals, Leicestershire, UK..

5 **Example 9: Determination of In Vitro Cytotoxicity**

K562 human chronic myeloid leukaemia cells were maintained in RPM1 1640 medium supplemented with 10% fetal calf serum and 2 mM glutamine at 37°C in a humidified atmosphere containing 5% CO₂ and were incubated with a specified dose of drug for 1 hour or 96 hours at 37°C in the dark. The incubation was terminated by centrifugation (5 min, 300 g) and the cells were washed once with drug-free medium. Following the appropriate drug treatment, the cells were transferred to 96-well microtiter plates (10⁴ cells per well, 8 wells per sample). Plates were then kept in the dark at 37°C in a humidified atmosphere containing 5% CO₂. The assay is based on the ability of viable cells to reduce a yellow soluble tetrazolium salt, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT, Aldrich-Sigma), to an insoluble purple formazan precipitate. Following incubation of the plates for 4 days (to allow control cells to increase in number by approximately 10 fold), 20 µL of MTT solution (5 mg/mL in phosphate-buffered saline) was added to each well and the plates further incubated for 5 h. The plates were then centrifuged for 5 min at 300 g and the bulk of the medium pipetted from the cell pellet leaving 10-20 µL per well. DMSO (200 µL) was added to each well and the samples agitated to ensure complete mixing. The optical density was then read at a wavelength of 550 nm on a Titertek Multiscan ELISA plate reader, and a dose-response curve was constructed. For each curve, an IC₅₀ value was read as the dose required to reduce the final optical density to 50% of the control value.

Results

The following compounds showed an IC₅₀ of less than 1µM after a 96 hour incubation period: 30a, 30b, 30c, 30d, 30f, 30g, 30h, 30i,

30j, 30k, 30l, 30m, 30n, 30o, 30p, 30q, 30r, 30s, 30t, 30u, 30v,
30w, 30x, 30y, 30z, 30aa, 30ac, 30ad, 30ae, 30af, 30ag, 30ah, 30ai,
30aj, 30al.


- 5 The following compounds showed an IC_{50} of less than 10nM after a 96 hour incubation period: 30a, 30b, 30c, 30g, 30i, 30n, 30p, 30q, 30ai.

Compound	IC_{50}^b (μ M)
ZC-204	3.80 ± 0.44
ZC-207	0.0053 ± 0.0049
ZC-209	<0.01
ZC-211	<0.01

^b 1 hour incubation

CLAIMS

1. A compound of formula I:



- 5 and salts, solvates and chemically protected forms thereof,
wherein:
R⁶ and R⁹ are independently selected from H, R, OH, OR, SH, SR, NH₂,
NHR, NRR', nitro, Me₃Sn and halo;
R and R' are independently selected from optionally substituted
10 C₁₋₁₂ alkyl, C₃₋₂₀ heterocyclyl and C₅₋₂₀ aryl groups;
R⁷ and R⁸ are independently selected from H, R, OH, OR, SH, SR, NH₂,
NHR, NRR', nitro, Me₃Sn and halo,
or the compound is a dimer with each monomer being of formula (I),
where the R⁷ groups or R⁸ groups of each monomers form together a
15 dimer bridge having the formula -X-R"-X- linking the monomers,
where R" is a C₃₋₁₂ alkylene group, which chain may be interrupted
by one or more heteroatoms and/or aromatic rings, and each X is
independently selected from O, S, or NH;
or any pair of adjacent groups from R⁶ to R⁹ together form a group
20 -O-(CH₂)_p-O-, where p is 1 or 2;
R¹⁰ is a carbamate-based nitrogen protecting group;
R¹¹ is an oxygen protecting group; and
R² is a labile leaving group.
- 25 2. A compound according to claim 1, wherein R⁹ is H.
3. A compound according to either claim 1 or claim 2, wherein R⁶
is selected from H, OH, OR, SH, NH₂, nitro and halo.
- 30 4. A compound according to any one of the preceding claims,
wherein R¹⁰ is Troc.

5. A compound according to any one of the preceding claims,
wherein R^{11} is a silyl oxygen protecting group or THP.

6. A compound according to any one of the preceding claims,
5 wherein R^2 is triflate.


7. A compound according to any one of the preceding claims,
wherein R^7 and R^8 are independently selected from H, OH, OR, SH,
10 NH_2 , NHR, NRR' and halo.

8. A compound according to any one of claims 1 to 6, wherein the
compound is a dimer with each monomer being of formula (I), where
the R^7 groups or R^8 groups of each monomer form together a dimer
bridge having the formula $-O-(CH_2)_n-O-$ linking the monomers, where n
15 is from 3 to 12.

9. A compound according to claim 8, wherein n is from 3 to 7.

10. A compound according to either claim 8 or claim 9, wherein
20 the substituents R^8 join to form the dimer bridge.

11. A compound of formula III:



25 and salts, solvates, chemically protected forms and prodrugs
thereof, wherein:

R^6 and R^9 are independently selected from H, R, OH, OR, SH, SR, NH_2 ,
NHR, NRR', nitro, Me_3Sn and halo;

R and R' are independently selected from optionally substituted
 C_{1-12} alkyl, C_{3-20} heterocyclyl and C_{5-20} aryl groups;

30 R^7 and R^8 are independently selected from H, R, OH, OR, SH, SR, NH_2 ,
NHR, NRR', nitro, Me_3Sn and halo,

- or the compound is a dimer with each monomer being of formula (I), where the R⁷ groups or R⁸ groups of each monomers form together a dimer bridge having the formula -X-R"-X- linking the monomers, where R" is a C₃₋₁₂ alkylene group, which chain may be interrupted by one or more heteroatoms and/or aromatic rings, and each X is independently selected from O, S, or NH;
or any pair of adjacent groups from R⁶ to R⁹ together form a group -O-(CH₂)_p-O-, where p is 1 or 2;
R¹⁰ is a carbamate-based nitrogen protecting group; and
R¹⁶ is either O-R¹¹, wherein R¹¹ is an oxygen protecting group, or O-R¹¹ is OH; or
R¹⁰ and R¹⁶ together form a double bond between N10 and C11;
R¹⁵ is R;
and wherein,
when R⁷ and R⁸ are OMe, R⁶ and R⁹ are H, and where R¹⁰ and R¹⁶ together form a double bond between N10 and C11, R¹⁵ is not phenyl, 4-methylphenyl, 2-methylphenyl, 4-ethylphenyl, 2,6-dimethylphenyl, 4-methoxyphenyl, 4-tert-butylphenyl, 4-fluorophenyl, 4-chlorophenyl, 2-naphthyl or 2-thiophenyl.
12. A compound according to claim 11, wherein when R⁷ and R⁸ are OMe, R⁶ and R⁹ are H, and R¹⁵ is R, R is selected from the group 3-methoxyphenyl, 4-biphenyl, 4-phenoxyphenyl, 3,4-methylenedioxyphenyl, trans-2-(4-methylphenyl)vinyl, trans-propenyl, 4-dimethylaminophenyl, 4-methylthiophenyl, 4-vinylphenyl, 3,4-dichlorophenyl, 4-trifluoromethylphenyl, 4-isopropylphenyl, 4-cyanophenyl, 3-pyridinyl, 4-pyridinyl, 4-formylphenyl, 4-carboxylphenyl, 2,6-dimethoxyphenyl, 4-acetanilide, 4-aminophenyl, 1-naphthyl, 5-indole, 3-aminophenyl, 2,6-difluorophenyl, 1-pyrenyl, 4-hydroxyphenyl and trans-hexenyl.
13. A compound according to either claim 11 or claim 12, wherein when R⁷ and R⁸ are OMe, R⁶ and R⁹ are H, and R¹⁵ is R, R is selected from a C₃₋₂₀ heterocyclyl group having a nitrogen ring atom, C₅₋₂₀ aryl group having a nitrogen-containing substituent, or a C₅₋₂₀

heteroaryl group having a nitrogen ring atom or a nitrogen-containing substituent.

14. A compound according to claim 11, wherein the compound is a dimer with each monomer being of formula (I), where the R⁷ groups or R⁸ groups of each monomer form together a dimer bridge having the formula -O-(CH₂)_n-O- linking the monomers, where n is from 3 to 12.

15. A compound according to claim 14, wherein n is from 3 to 7.

16. A compound according to either claim 14 or claim 15, wherein the substituents R⁸ join to form the dimer bridge.

17. A compound according to any one of claims 14 to 16, wherein R¹⁵ is selected from:

- (i) optionally substituted C₅₋₂₀ aryl groups;
- (ii) substituted C₂ alkyl groups; and
- (iii) optionally substituted C₃₋₇ alkyl groups.

18. A compound according to any one of claims 11 to 17, wherein R¹⁰ and R¹⁶ together form a double bond between N10 and C11.

19. A compound according to any one of claims 11 to 18, wherein R⁹ is H.

20. A compound according to any one of claims 11 to 19, wherein R⁷ and R⁸ are independently selected from H, OH, OR, SH, NH₂, NHR, NRR', and halo.


21. A compound according to any one of claims 11 to 20 for use in a method of therapy.

22. A pharmaceutical composition containing a compound of any one of claims 11 to 20, and a pharmaceutically acceptable carrier or diluent.


5 23. Use of a compound according to any one of claims 11 to 20 in the manufacture of a medicament for treating a proliferative disease.

10 24. A method of treatment of a proliferative disease, comprising administering to a subject in need of treatment a therapeutically-effective amount of a compound of any one of claims 11 to 20.

25. A method of synthesising a compound of formula I:



15 from a compound of formula IIa:



wherein:

R⁶ and R⁹ are independently selected from H, R, OH, OR, SH, SR, NH₂, NHR, NRR', nitro, Me₃Sn and halo;

20 R and R' are independently selected from optionally substituted C₁₋₁₂ alkyl, C₃₋₂₀ heterocyclyl and C₅₋₂₀ aryl groups;

R⁷ and R⁸ are independently selected from H, R, OH, OR, SH, SR, NH₂, NHR, NRR', nitro, Me₃Sn and halo,

or the compound is a dimer with each monomer being of formula (I),

25 where the R⁷ groups or R⁸ groups of each monomers form together a dimer bridge having the formula -X-R''-X- linking the monomers, where R'' is a C₃₋₁₂ alkylene group, which chain may be interrupted

by one or more heteroatoms and/or aromatic rings, and each X is independently selected from O, S, or NH;
 or any pair of adjacent groups from R⁶ to R⁹ together form a group -O-(CH₂)_p-O-, where p is 1 or 2;


5 R¹⁰ is a carbamate-based nitrogen protecting group;

R¹¹ is an oxygen protecting group;

R² is a labile leaving group; and

R¹² and R¹³ together form =O.

10 26. A method according to claim 25, wherein the compound of formula IIa is synthesised from a compound of formula IIb:



wherein said compound of formula IIb has R⁶, R⁷, R⁸, R⁹, R¹⁰ and R¹¹ defined according to claim 25, and for said compound of formula IIb
 15 R¹² is O-R¹⁴, and R¹³ is H; and
 R¹⁴ is an oxygen protecting group orthogonal to R¹¹.


27. A method according to claim 26, wherein the compound of formula IIa is synthesised using an oxidation reaction performed
 20 under Swern conditions, or a method involving the TPAP or the Dess Martin reagents.

28. A method according to any one of claims 25 to 27, wherein when R² in the compound of formula I is -OSO₂CH₃, -OSO₂(C_nF_{2n+1}) where
 25 n = 0, 1 or 4, or -OSO₂R⁵ where R⁵ is an optionally substituted phenyl group, then said compound of formula I is synthesised by using a treatment step with the appropriate R² anhydride.


29. A method according to any one of claims 25 to 27, wherein
 30 when R² in the compound of formula I is -I or -Br, then said compound of formula I is synthesised by using a reaction step involving hydrazine and iodine or bromine.

30. A method according to any one of claims 25 to 27, wherein
 when R^2 in the compound of formula I is $-Cl$, then said compound of
 formula I is synthesised by using a reaction step involving
 5 phosphorous oxychloride.

31. A method of synthesising a compound of formula III:



from a compound of formula I:



10

wherein

R^6 and R^9 are independently selected from H, R, OH, OR, SH, SR, NH_2 ,
 NHR, NRR' , nitro, Me_3Sn and halo;

R and R' are independently selected from optionally substituted
 15 C_{1-12} alkyl, C_{3-20} heterocyclyl and C_{5-20} aryl groups;

R^7 and R^8 are independently selected from H, R, OH, OR, SH, SR, NH_2 ,
 NHR, NRR' , nitro, Me_3Sn and halo,

or the compound is a dimer with each monomer being of formula (I),
 where the R^7 groups or R^8 groups of each monomers form together a
 dimer bridge having the formula $-X-R''-X-$ linking the monomers,
 where R'' is a C_{3-12} alkylene group, which chain may be interrupted
 by one or more heteroatoms and/or aromatic rings, and each X is
 20 independently selected from O, S, or NH;

or any pair of adjacent groups from R^6 to R^9 together form a group
 25 $-O-(CH_2)_p-O-$, where p is 1 or 2;

R^{10} is a carbamate-based nitrogen protecting group;

R^2 is a labile leaving group;

R¹⁶ is either O-R¹¹, where R¹¹ is an oxygen protecting group, or OH, or R¹⁰ and R¹⁶ together form a double bond between N10 and C11; and R¹⁵ is R.

5 32. A method according to claim 31, wherein the synthesis of said compound of formula III uses a palladium catalysed coupling step.

33. A method according to claim 32, wherein the palladium catalyst is Pd(PPh₃)₄, Pd(OCOCH₃)₂, PdCl₂ or Pd(dba)₃.

10

34. A method according to either claim 32 or claim 33, wherein the coupling reaction is performed under microwave conditions.

15 35. A method according to any one of claims 32 to 34, wherein the palladium catalyst is solid supported.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB2005/000768

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 C07D519/00 A61K31/5517 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEN ET AL: "A novel approach to the synthesis of cytotoxic C2?C3 unsaturated pyrrolo'2,1-c!' and !benzodiazepines (PBDs) with conjugated acrylyl C2-substituents" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, vol. 14, no. 6, 27 February 2004 (2004-02-27), pages 1547-1549, XP002329273 page 1548; figure 1	1-35
X	KANG ET AL: "Synthesis of a novel C2-aryl substituted 1,2-unsaturated pyrrolobenzodiazepine" CHEMICAL COMMUNICATIONS, vol. 14, 11 June 2003 (2003-06-11), pages 1688-1689, XP002329274 page 1688	11-24

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

24 May 2005

Date of mailing of the international search report

20/07/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Cortés, J

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB2005/000768

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GREGSON ET AL: "Synthesis of the first examples of A-C8/C-C2 amide-Linked pyrrolo'2,1-c!'1,4!benzodiazepine dimers" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, vol. 13, no. 14, 31 March 2003 (2003-03-31), pages 2277-2280, XP002329063 scheme 3 page 2279	11-24
X	COOPER ET AL: "Synthesis of novel C2-aryl pyrrolobenzodiazepines (PBDs) as potential antitumour agents" CHEMICAL COMMUNICATIONS, vol. 16, 5 July 2002 (2002-07-05), pages 1764-1765, XP002329275 page 1764	11-24
X	GREGSON ET AL: "Synthesis of the first example of a c2-C3/C2?-C3?-endo unsaturated pyrrolo'2,1-c!'1,4!benzodiazepine dimer" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, vol. 11, no. 21, 27 August 2001 (2001-08-27), pages 2859-2862, XP002329276 page 2861	11-24
X	GREGSON ET AL: "Effect of C2/C3-endo unsaturation on the cytotoxicity and dna-binding reactivity of pyrrolo'2,1-c!'1,4!benzodiazepines" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, vol. 10, no. 16, 14 April 2000 (2000-04-14), pages 1849-1852, XP004216014 figure 1, schemes 1-3 page 1850 - page 1851	11-24
X	WO 00/12508 A (THE UNIVERSITY OF PORTSMOUTH HIGHER EDUCATION (GB)) 9 March 2000 (2000-03-09) cited in the application the whole document	11-24
X	FUKUYAMA ET AL: "Total synthesis of (+)-porothramycin B" TETRAHEDRON LETTERS, vol. 34, no. 16, 16 April 1993 (1993-04-16), pages 2577-2580, XP002135999 page 2577 page 2578	11-24
	-/-	

INTERNATIONAL SEARCH REPORT

Inter ☐ if Application No
PCT/GB2005/000768

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 2004/043963 A (SPIROGEN (GB)) 27 May 2004 (2004-05-27) the whole document	11-24
P,X	TIBERGHIE ET AL: "Application of the Stille coupling reaction to the synthesis of C2-substituted endo-exo unsaturated pyrrolo'2,1-c!'1,4!benzodiazepines (PBDs)" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, vol. 14, no. 20, 1 September 2004 (2004-09-01), pages 5041-5044, XP002329277 the whole document	11-24

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB2005/000768

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: --
because they relate to subject matter not required to be searched by this Authority, namely:
Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/GB2005/000768

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0012508	A	09-03-2000	AT 240334 T	15-05-2003
			AT 294803 T	15-05-2005
			AU 757510 B2	20-02-2003
			AU 5635199 A	21-03-2000
			CA 2341471 A1	09-03-2000
			DE 69907977 D1	18-06-2003
			DE 69907977 T2	22-07-2004
			DE 69925133 D1	09-06-2005
			DK 1193270 T3	15-09-2003
			EP 1193270 A2	03-04-2002
			EP 1413582 A1	28-04-2004
			EP 1109812 A2	27-06-2001
			ES 2199200 T3	16-02-2004
			WO 0012508 A2	09-03-2000
			JP 2002525285 T	13-08-2002
			NZ 510493 A	28-11-2003
			PT 1193270 T	31-10-2003
			US 2003120069 A1	26-06-2003
WO 2004043963	A	27-05-2004	AU 2003282251 A1	03-06-2004
			WO 2004043963 A1	27-05-2004

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)